

S1 Table. Sociodemographic, clinical, immunological, and virological profile of the HIV-infected children at diagnosis, by origin of the children

	Spanish N=406	Sub-Saharan Africa N=82	Latin America N=27	Other regions N=17
Calendar Period, N (%)				
1997-2000	191 (47)	6 (7.3)	3 (11.1)	3 (17.6)
2001-2005	118 (29.1)	21 (25.6)	7 (25.9)	3 (17.6)
2006-2010	73 (18)	38 (46.3)	12 (44.4)	7 (41.2)
2011-2015	24 (5.9)	17 (20.7)	5 (18.5)	4 (23.5)
Sex, N (%)				
Males	187 (46.1)	42 (51.2)	13 (48.1)	9 (52.9)
Females	219 (53.9)	40 (48.8)	14 (51.9)	8 (47.1)
HIV transmission route, N (%)				
Perinatal	387 (95.3)	61 (74.4)	14 (51.9)	14 (82.3)
Transfusional	3 (0.7)	8 (9.8)	3 (11.1)	0 (0)
Sexual	6 (1.5)	0 (0)	8 (29.6)	2 (11.8)
Unknown (mother not HIV infected)	10 (2.5)	2 (2.4)	0 (0)	0 (0)
Unknown (HIV mother status not known)	0 (0)	11 (13.4)	2 (7.4)	1 (5.9)
CDC stage, N (%)	N=402	N=81	N=27	N=17
N-A	267 (66.4)	37 (45.7)	16 (59.3)	8 (47.1)
B	78 (19.4)	21 (25.9)	4 (14.8)	6 (35.3)
C	57 (14.2)	23 (28.4)	7 (25.9)	3 (17.6)
Coinfections, N (%)				
HCV	12 (3)	1 (1.2)	0 (0)	0 (0)
HBV	0 (0)	8 (9.8)	0 (0)	1 (5.9)
%CD4, Median (IQR)	N=346	N=78	N=26	N=15
	28 (15.2-40)	17.5 (9.5-24.8)	15.5 (5.2-23.8)	17 (10.5-25.5)
CD4/mm³, Median (IQR)	N=348	N=78	N=26	N=15
	1176 (465-2185)	609 (237-1143)	389 (68-703)	379 (239-747)
Log Viral Load, Median (IQR)	N=353	N=77	N=26	N=15
	5.2 (4.6-5.9)	5.1 (4.4-5.6)	5.1 (4.7-5.7)	5 (4.4-5.6)

Other regions: Eastern Europe (N=6), North Africa (N=5), Western Europe (N=3), Asia (N=3)