

S2 Table. SocioDemographic, clinical, immunological, and virological profile at diagnosis of perinatally HIV-infected children born in Spain and their mothers, Spanish HIV-infected children at diagnosis, by origin of the mothers

	Spanish N=299	Sub-Saharan Africa N=39	Latin America N=19	Other regions N=16
<b>Calendar Period, N (%)</b>				
1997-2000	163 (54.5)	5 (12.8)	5 (26.3)	1 (6.3)
2001-2005	95 (31.8)	11 (28.2)	5 (26.3)	4 (25)
2006-2010	32 (10.7)	17 (43.6)	5 (26.3)	11 (68.7)
2011-2015	9 (3)	6 (15.4)	4 (21.1)	0 (0)
<b>Period of HIV diagnosis for mothers, N (%)</b>				
Until childbirth	140 (46.8)	21 (53.8)	10 (52.6)	11 (68.7)
After childbirth	159 (53.2)	18 (46.2)	9 (47.4)	5 (31.3)
<b>Mode of HIV transmission of mothers, N (%)</b>	N=281	N=38	N=18	N=15
IDU	146 (52)	1 (2.6)	2 (11.1)	1 (6.7)
No history of IDU	135 (48)	37 (97.4)	16 (88.9)	14 (93.3)
<b>Sex of children, N (%)</b>				
Male	139 (46.5)	15 (38.5)	4 (21.1)	10 (50)
Female	160 (53.5)	24 (61.5)	15 (78.9)	6 (50)
<b>Age of children, N (%)</b>				
<1 year	188 (62.9)	27 (69.2)	12 (63.2)	12 (83.3)
1-5 years	76 (25.4)	11 (28.2)	6 (31.6)	4 (16.7)
>5 years	35 (11.7)	1 (2.6)	1 (5.3)	0 (0)
<b>Coinfections in children, N (%)</b>				
HCV	10 (3.3)	0 (0)	1 (5.3)	0 (0)
HBV	0 (0)	0 (0)	0 (0)	0 (0)
<b>CDC stage of children, N (%)</b>	N=295	N=39	N=19	N=16
N-A	190 (64.4)	32 (82.1)	13 (68.4)	14 (87.5)
B	60 (20.3)	3 (7.7)	3 (15.8)	1 (6.3)
C	45 (15.3)	4 (10.3)	3 (15.8)	1 (6.3)
<b>%CD4, Median (IQR)</b>	N=247	N=35	N=19	N=14
	27 (15-38.7)	31 (21.9-44.3)	38.3 (22-46.5)	36 (30.4-44)
<b>CD4/mm<sup>3</sup>, Median (IQR)</b>	N=249	N=35	N=19	N=14

<b>Log Viral Load, Median (IQR)</b>	1180 (465-2142)	1518 (915-2885)	1928 (638-2239)	2119 (972-3068)
	N=256	N=35	N=17	N=13
	5.4 (4.8-5.9)	4.8 (4-5.9)	5.2 (4.4-5.9)	5.7 (5.1-5.9)

Other regions: Eastern Europe (N=6), North Africa (N=6), [Western](#) Europe (N=4)