

Authors:

Begoña Vieites¹, María Ángeles López-García^{1,11}, María Dolores Martín Salvago², Cesar Luis Ramirez Tortosa², Ricardo Rezola³, Sancho Magdalena⁴, Laura López Vilaró⁵, Felip Vilardell Villellas⁶, Octavio Burgués⁷, Beatriz Fernández-Rodriguez⁸, Lina Alfaro Galán⁹, Vicente Peg^{10,11}

Title:

"Predictive and prognostic value of total tumor load in sentinel lymph nodes in breast cancer patients after neoadjuvant treatment using one-step nucleic acid amplification: the NEOVATTL study"

Affiliations:

¹Department of Pathology - Hospital Universitario Virgen del Rocío (Sevilla, Spain)

²Department of Pathology - Hospital Universitario Materno-Infantil (Jaén, Spain)

³Department of Pathology - Onkologikoa Kutxa Fundazioa (Donostia, Spain)

⁴Department of Pathology - Hospital Universitario de Salamanca (Salamanca, Spain)

⁵Department of Pathology- Hospital de la Santa Creu i Sant Pau (Barcelona, Spain)

⁶Hospital Universitari Arnau de Vilanova (Lérida, Spain)

⁷Department of Pathology - Hospital Clínico Universitario de Valencia (Valencia, Spain)

⁸Department of Pathology - Complejo Hospitalario Universitario de Santiago (Santiago de Compostela, Spain)

⁹Department of Gynaecology and Obstetrics - Hospital Universitario Virgen del Rocío (Sevilla, Spain)

¹⁰Department of Pathology - Hospital Universitari Vall d'Hebron (Barcelona, Spain)

¹¹CIBERONC (Centro de Investigación Biomédica en Red de Cáncer) – Instituto de Salud Carlos III (Madrid, Spain)

Corresponding author: Begoña Vieites

Email: mb.vieites.sspa@juntadeandalucia.es

Online Resource 3

Supplementary Fig. S1 ROC curve based on the sensitivity and the specificity of TTL for the prediction of non-SNL involvement. ROC, receiver operator characteristic; SNL, sentinel lymph node; TTL, total tumor load

