

# First-in-Human Phase I Study of Iadademstat (ORY-1001): A First-in-Class Lysine-Specific Histone Demethylase 1A Inhibitor, in Relapsed or Refractory Acute Myeloid Leukemia

Olga Salameo, MD<sup>1</sup>; Pau Montesinos, MD<sup>2,3</sup>; Christophe Willekens, MD<sup>4</sup>; José Antonio Pérez-Simón, MD, PhD<sup>5,6</sup>; Arnaud Pigneux, MD, PhD<sup>7</sup>; Christian Récher, MD, PhD<sup>8</sup>; Rakesh Popat, MB, BS, PhD<sup>9</sup>; Cecilia Carpio, MD<sup>1</sup>; César Molinero, MD, PhD<sup>10</sup>; Cristina Mascaró, PhD<sup>10</sup>; Joaquim Vila<sup>10</sup>; M. Isabel Arévalo, PhD<sup>10</sup>; Tamara Maes, PhD<sup>10</sup>; Carlos Buesa, PhD<sup>10</sup>; Francesc Bosch, MD, PhD<sup>1</sup>; and Tim C. P. Somervaille, MBBS, PhD<sup>11,12</sup>

abstract

**PURPOSE** Iadademstat is a novel, highly potent, and selective inhibitor of LSD1 (KDM1A), with preclinical in vitro and in vivo antileukemic activity. This study aimed to determine safety and tolerability of iadademstat as monotherapy in patients with relapsed/refractory acute myeloid leukemia (R/R AML).

**METHODS** This phase I, nonrandomized, open-label, dose-escalation (DE), and extension-cohort (EC) trial included patients with R/R AML and evaluated the safety, pharmacokinetics (PK), pharmacodynamics (PD), and preliminary antileukemic activity of this orally bioavailable first-in-class lysine-specific demethylase 1 inhibitor.

**RESULTS** Twenty-seven patients were treated with iadademstat on days 1 to 5 (5-220  $\mu\text{g}/\text{m}^2/\text{d}$ ) of each week in 28-day cycles in a DE phase that resulted in a recommended dose of 140  $\mu\text{g}/\text{m}^2/\text{d}$  of iadademstat as a single agent. This dose was chosen to treat all patients ( $n = 14$ ) in an EC enriched with patients with MLL/KMT2A-rearranged AML. Most adverse events (AEs) were as expected in R/R AML and included myelosuppression and nonhematologic AEs, such as infections, asthenia, mucositis, and diarrhea. PK data demonstrated a dose-dependent increase in plasma exposure, and PD data confirmed a potent time- and exposure-dependent induction of differentiation biomarkers. Reductions in blood and bone marrow blast percentages were observed, together with induction of blast cell differentiation, in particular, in patients with MLL translocations. One complete remission with incomplete count recovery was observed in the DE arm.

**CONCLUSION** Iadademstat exhibits a good safety profile together with signs of clinical and biologic activity as a single agent in patients with R/R AML. A phase II trial of iadademstat in combination with azacitidine is ongoing (EudraCT No.: 2018-000482-36).

J Clin Oncol 38:4260-4273. © 2020 by American Society of Clinical Oncology

Creative Commons Attribution Non-Commercial No Derivatives 4.0 License 

ASSOCIATED CONTENT

Data Supplement Protocol

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on September 8, 2020 and published at [ascopubs.org/journal/jco](https://ascopubs.org/journal/jco) on October 14, 2020; DOI <https://doi.org/10.1200/JCO.19.03250>

## INTRODUCTION

Acute myeloid leukemia (AML) is a hematologic malignancy characterized by a myeloid lineage differentiation block.<sup>1-3</sup> It remains, for the most part, an incurable disease, especially in the elderly, and new approaches to treatment are required, including those that promote differentiation.<sup>4-10</sup> Epigenetic dysfunction has a central role in AML pathology, as evidenced by recurrent mutations in transcription factors and epigenetic regulators.<sup>11-14</sup> Certain regulators are under evaluation as therapeutic targets, including lysine-specific demethylase 1 (LSD1), which serves a dual role in hematopoiesis. It exhibits demethylase activity versus monomethylated and dimethylated lysine residues on histone tails<sup>15</sup> as well as scaffolding activity,

which facilitates recruitment of histone deacetylase to sites on chromatin where SNAG domain transcription factors such as GFI1 and GFI1B are bound.<sup>16</sup> LSD1 is highly expressed in hematopoietic stem cells and myeloblasts, and is necessary for proliferation and terminal differentiation during normal hematopoiesis.<sup>17</sup> Preclinical studies have revealed that LSD1 sustains the differentiation block in certain molecular subtypes of AML, in particular, MLL-translocated AML, and is required for leukemic stem cell potential.<sup>18-22</sup> Targeting LSD1 in AML may serve to promote differentiation of leukemic blasts.

Iadademstat (ORY-1001) is a highly selective and potent covalent inhibitor of LSD1, which induces differentiation of AML cells in vitro at low concentrations

## CONTEXT

### Key Objectives

To evaluate the safety, pharmacokinetics, pharmacodynamics and preliminary antileukemic activity of iadademstat in acute myeloid leukemia.

### Knowledge Generated

Iadademstat exhibits a good safety profile together with signs of clinical and biologic activity as a single agent in patients with relapsed or refractory AML.

### Relevance

Current treatment options in AML fail to cure the majority of patients, in particular those not fit for intensive chemotherapy, and novel therapies are required. Ongoing studies are investigating the combinatorial use of iadademstat with azacitidine to further delineate its activity in AML.

(< 1 nM) and compromises leukemic stem cell capacity in preclinical models of AML. Iadademstat induces a monocyte/macrophage differentiation gene signature in AML cell lines, and induction of differentiation biomarkers correlates with reduction of tumor growth in rodent leukemia xenografts. Iadademstat has excellent oral bioavailability, excellent target exposure, and promising antitumor activity in vivo.<sup>23</sup> We report a first-in-human dose-escalation (DE) and extension-cohort (EC) phase I study with iadademstat in patients with refractory or relapsed (R/R) acute leukemia (EudraCT No.: 2013-002447-29). The primary objective was to assess safety and tolerability of iadademstat; secondary objectives were to study pharmacokinetics (PK), pharmacodynamics (PD), and efficacy.

## METHODS

### Participant Selection

Patients with relapsed or refractory acute leukemia (excluding acute promyelocytic leukemia; age  $\geq$  16 years) not deemed suitable for standard therapies, with Eastern Cooperative Oncology Group performance status  $\leq$  2 and without an unstable or uncontrolled concurrent severe medical condition, were eligible. Prior acute leukemia treatment should have stopped at least 14 days before the first dose of iadademstat. Hydroxycarbamide was allowed until 12 hours before the first dose of iadademstat and then after the fifth day (Data Supplement, online only). Patients with *MLL*-rearranged AML or acute erythroblastic leukemia were selected for the EC.

### Study Design

The protocol was approved by the institutional review board or independent ethics committee at each participating center and by regulatory authorities, in accordance with the Declaration of Helsinki, the International Conference on Harmonization–Good Clinical Practice, and local laws. Investigators obtained informed consent from each participant before performing any study-specific procedures. Data were anonymized to protect patient identities.

DE was performed in a traditional 3 + 3 design (Data Supplement),<sup>24,25</sup> with dosing administered until disease progression, death, consent withdrawal, or adverse events (AEs) that did not improve by standard of care (Data Supplement). The starting dose was based on preclinical toxicology studies (Data Supplement). The escalation doses were 5, 15, 30, 45, 60, 80, 140, and 220  $\mu\text{g}/\text{m}^2/\text{d}$  in cohorts 1 to 8, the latter included after an amendment to add an additional dose level (220  $\mu\text{g}/\text{m}^2/\text{d}$ ). Iadademstat was administered orally after a minimum of 2 hours of fasting using a precharged syringe in a 28-day cycle (5 days on/2 days off  $\times$  4 weeks). Details regarding maximum tolerated dose assessment, study monitoring, safety analysis, sampling, PK/PD analysis, bone marrow (BM) biomarker analysis, response analysis, and statistical assessments are available in the Data Supplement.

## RESULTS

### Patient Characteristics

Between February 2014 and April 2015, 27 patients with R/R acute leukemia were enrolled in the DE phase (1 with ALL in cohort 1; 26 with AML). Two patients failed screening. Between September 2015 and May 2016, 14 patients were enrolled in the EC (140  $\mu\text{g}/\text{m}^2/\text{d}$ ), which, guided by preclinical data, was restricted to patients with *MLL* gene rearrangements or erythroleukemia. Median age was 67 years; 15 (42%) of 36 patients with karyotypes available exhibited adverse risk cytogenetics,<sup>26</sup> and 7 (17%) of 41 patients were in second or third relapse. The median time since initial diagnosis to study enrollment was 9.8 months and since last treatment was 1.3 months. Of 41 patients, 30 completed cycle 1 (C1; Table 1; Data Supplement).

### Safety and Tolerability

All patients experienced treatment-emergent AEs; in total, 497 were reported (Data Supplement). The most frequent nonhematologic AEs were infection, GI symptoms, hemorrhagic manifestations, asthenia, musculoskeletal

**TABLE 1.** Summary of Patient Characteristics at Screening

Characteristics	Dose-Escalation Cohort	Extension Cohort	All Patients
No. of patients	27	14	41
Median age, years (range),	69 (40-81)	63 (30-78)	67 (30-81)
Sex (F/M), %	30/70	36/64	32/68
ECOG performance status, No.			
0	4	2	6
1	16	11	27
2	5	1	6
NA	2	0	2
Median WBC count $\times 10^9/L$ , (range)	4.0 (0.6-104.9)	2.5 (0.2-26.3)	2.9 (0.2-104.9)
Median platelet count $\times 10^9/L$ , (range)	31 (4-335)	19 (4-266)	21 (4-335)
Median Hgb, g/L (range)	89 (67-122)	86 (74-143)	88 (67-143)
Median PB blast, % (range)	42 (0-92)	31 (0-95)	39 (0-95)
Median BM blast, % (range),	49 (0-90)	32 (2-97)	47 (0-97)
Cytogenetic risk, <sup>26</sup> No.			
Favorable	1	0	1
Intermediate	12	8	20
Adverse	10	5	15
NA	4	1	5
Molecular abnormalities, No.			
<i>MLL</i> rearrangements	0	6	6
<i>MLL</i> , other abnormalities	2	4	6
FLT3-ITD mutations	3	0	3
NPM1 mutations	2	0	2
WT1	8	2	10
TP53 mutations	1	1	2
NA	14	4	18
WHO diagnosis, <sup>1-4</sup> No.			
AML with recurrent genetic abnormalities	10	4	14
AML with multilineage dysplasia	6	5	11
AML, therapy related	2	2	4
AML, not otherwise specified	5	3	8
NA	4	0	4
FAB classification, <sup>1-4</sup> No.			
M0	3	0	3
M2	4	3	7
M4	0	4	4
M5	1	3	4
M6	0	4	4
M7	2	0	2
NA	17	0	17
No. of relapses			
0 (refractory)	8	5	13
1	14	7	21
$\geq 2$	5	2	7
Median time from AML diagnosis months, (range)	9.8 (2-41)	9.9 (4-36)	9.8 (2-41)

Abbreviations: AML, acute myeloid leukemia; BM, bone marrow; ECOG, Eastern Cooperative Oncology Group; Hgb, hemoglobin; NA, not available; PB, peripheral blood.

**TABLE 2.** Grade 3 and 4 AEs and SAEs

AE Description	Total Grade 3 and 4 AEs		Grade 3 and 4 AEs Considered Possibly, Probably, or Definitely Related to ladademstat	
	DE (n = 27)	EC (n = 14)	DE (n = 27)	EC (n = 14)
General				
Asthenia	2	4	—	2
Anorexia	—	2	—	2
Mucositis	1	—	—	—
Edema	1	1	—	—
Infection				
Fever of unknown origin or neutropenic fever	10	11	1	3
Septic shock or sepsis	2	3	—	—
Cutaneous/subcutaneous infection	3	2	—	—
Respiratory infection or pneumonia	5	2	—	1
Tonsillitis	—	1	—	—
Hemorrhage				
Cutaneous	2	—	—	—
Mucosal	1	—	—	—
Intracranial	1	—	—	—
Respiratory (other than mentioned)				
Respiratory failure or distress	1	—	—	—
Pleural effusion	2	—	—	—
GI (other than mentioned)				
Abdominal pain	1	—	—	—
Neurologic				
Seizure or vasovagal episode	2	—	—	—
Hematologic				
Thrombocytopenia	8	3	3	2
Leukocytosis	3	1	—	—
Leukopenia	1	—	—	—
Neutropenia	3	1	2	—
Anemia	1	1	—	—
Musculoskeletal disorders				
Musculoskeletal pain or discomfort	—	1	—	—
Metabolism disorders				
Hyperglycemia	1	—	—	—
Hypokalemia	7	—	—	—
Cardiac				
Supraventricular tachycardia	—	2	—	—
Hypotension	2	1	—	—
Urinary tract disorders				
Renal impairment	—	1	—	—

(continued on following page)

**TABLE 2.** Grade 3 and 4 AEs and SAEs (continued)

AE Description	Total Grade 3 and 4 AEs		Grade 3 and 4 AEs Considered Possibly, Probably, or Definitely Related to ladademstat	
	DE (n = 27)	EC (n = 14)	DE (n = 27)	EC (n = 14)
Other				
Graft versus host disease	1	—	—	—
Infusion reaction	1	—	—	—
Total	62	37	6	10

SAE Description	Total SAEs		SAEs Considered Possibly, Probably, or Definitely Related to ladademstat	
	DE (n = 27)	EC (n = 14)	DE (n = 27)	EC (n = 14)
General				
Asthenia	—	1	—	1
Infection				
Fever of unknown origin or neutropenic fever	7	9	1	3
Septic shock or sepsis	3	2	—	—
Cutaneous/subcutaneous	1	2	—	1
Respiratory infection or pneumonia	10	3	1	1
Sinusitis	1	—	—	—
Bleeding				
Intracranial hemorrhage	3	—	—	—
Respiratory (other than mentioned)				
Respiratory failure or distress	2	—	—	—
GI (other than mentioned)				
Diarrhea	—	1	—	—
Neurologic				
Depressed level of consciousness	1	—	—	—
Hematologic				
Thrombocytopenia	—	1	—	1
Leukocytosis	—	3	—	—
Differentiation syndrome	—	2	—	2
Disease progression	1	5	—	—
Musculoskeletal				
Musculoskeletal pain or discomfort	—	1	—	—
Cardiac				
Pericarditis	—	1	—	—
Supraventricular tachycardia	—	2	—	—
Cardiac failure	1	—	—	—
Hypotension	—	1	—	—
Other				
Graft versus host disease (liver)	1	—	—	—
Infusion reaction	1	—	—	—
Total	32	34	2	9

NOTE. AEs that evolved from a lower to a higher grade or vice versa (n = 21) are only counted once, at the highest grade.  
Abbreviations: AE, adverse event; DE, dose escalation; EC, extension cohort; SAE, serious adverse event.

pain, mucositis, edema, skin rash, and anorexia. Of the AEs, 99 were grade 3 or 4; 66 were reported as serious AEs (SAEs), mainly infections (Table 2; Data Supplement).

Sixty-six AEs were considered related to iadademstat (Data Supplement), 10 certainly, 21 probably, and 35 possibly, according to investigator assessment. Of these, 16 were grade 3 or 4, and 11 were SAEs (Table 2; Data Supplement). SAEs included neutropenic fever or fever of unknown origin (n = 4), cellulitis (n = 1), pneumonia or respiratory infection (n = 2), thrombocytopenia (n = 1), asthenia (n = 1), and differentiation syndrome (n = 2).

During the study, there were 27 recorded grade 5 AEs (where the AE was considered by the investigator to have contributed to death; Table 2; Data Supplement), and 25 patients died (DE, n = 16; EC, n = 9). The recorded causes of death were as expected for patients with R/R acute leukemia: AML (n = 12), lung infection or respiratory failure (n = 9), septic shock (n = 1), sinusitis (n = 1), and intracranial hemorrhage with heart failure (n = 1). Three grade 5 AEs were considered possibly related to iadademstat: pneumonia in a patient in cohort VIII and episodes of cellulitis and differentiation syndrome in an EC patient whose recorded cause of death was AML. One EC patient died as a result of complications arising from a differentiation syndrome (n = 1), considered certainly related to iadademstat.

### Dose-Limiting Toxicity

In cohort VIII (220  $\mu\text{g}/\text{m}^2/\text{d}$ ), two SAEs were considered possibly related to iadademstat—pneumonia (patient 24; grade 5) and an episode of febrile neutropenia (patient 27; grade 3), which did not meet the strict protocol criteria for a dose-limiting toxicity (Data Supplement). However, in view of evidence in cohort VII (140  $\mu\text{g}/\text{m}^2/\text{d}$ ) of maximal biomarker induction within 24 hours of treatment, despite subsequent accumulation of plasma iadademstat from D1-D5 (Fig 1A and 1B; eg, patient 22), and evidence of potent hematopoietic target engagement (induction of grade 4 thrombocytopenia by day 15-17 of treatment in two patients; Figs 1C, and 1D), the Safety Monitoring Committee took the pragmatic view to establish the maximum tolerated dose as 220  $\mu\text{g}/\text{m}^2/\text{d}$ . The EC dose was set at 140  $\mu\text{g}/\text{m}^2/\text{d}$ . The effects of iadademstat on platelet levels across cohorts are shown in Fig 1C, and platelet dynamics for a cohort VII patient with high baseline platelet levels are shown Fig 1D. In the EC, one patient required a 25% dose reduction at cycle 1 day 15 (C1D15) after grade 4 thrombocytopenia and transient nonspecific deterioration in general health (patient 32).

### PK, PD, and Biomarkers

At 5 and 15  $\mu\text{g}/\text{m}^2/\text{d}$ , plasma concentrations were typically below the lower limit of quantification. At higher doses, concentrations increased in an approximately linear manner, with a tendency for overproportional exposure at doses > 80  $\mu\text{g}/\text{m}^2/\text{d}$ .  $T_{\text{max}}$  was generally observed 4-8


hours postdose. Compound accumulation was observed after repeated dosing, with an average accumulation ratio of approximately 3-6. The volume of distribution for iadademstat was approximately 200 times total body water, and the half-life was 40-100 hours. PK curves and parameters calculated for the EC (140  $\mu\text{g}/\text{m}^2/\text{d}$ ) are summarized in Figures 1A, 1B, and 1E.

As expected from preclinical data,<sup>23</sup> the PD biomarker response in the DE phase was heterogeneous. Nevertheless, selected biomarkers showed time and dose-dependent response profiles in individual patients. *PI16/CRISP9* was rapidly induced on day 1 (Fig 1B); maximal induction ( $-\Delta\Delta\text{Cp max}$ ) was achieved after multiple dosing in patient 11 (45  $\mu\text{g}/\text{m}^2/\text{d}$ ) and patient 18 (80  $\mu\text{g}/\text{m}^2/\text{d}$ ) but within 18 hours of the first dose in patient 22 (140  $\mu\text{g}/\text{m}^2/\text{d}$ ), reflecting saturation despite further increases in exposure over the following days. Remarkably, induction was sustained up to 1 week after the last administration. In the EC, patients with monocytic or monoblastic lineage leukemias (FAB-M4 or -M5 AML) exhibited a potent response of many biomarkers, with the highest increase observed for *VCAN*, *S100A12*, and *LY96*.

Biomarker response correlated with morphologic differentiation of blast cells in BM and/or peripheral blood (PB; Figs 2A-2C and Fig 3). A rapid and potent induction of *VCAN* and *S100A12* was observed in two patients who developed differentiation syndrome (patients 28 and 36). RNA sequencing of selected predose and post-treatment (C1D29) BM samples confirmed upregulation of *PROCR* and downregulation of erythroid biomarkers *GYPA*, *GYPB*, *HBA1*, and *HBB* in three patients with erythroleukemia (M6A) and confirmed the broad upregulation of biomarkers in PB in patient 29 (Data Supplement). Together, these surrogate biomarker data demonstrate target engagement of iadademstat in leukemic cells and support their utility as a tool to monitor response to LSD1 inhibition.

### Iadademstat Efficacy

In the DE phase, patient 16 achieved a complete remission with incomplete count recovery (CRi). This patient had FAB-M2 AML (Data Supplement), which had relapsed 6 months after a sibling donor allogeneic stem cell transplantation. At relapse, the patient received a single cycle of 60  $\mu\text{g}/\text{m}^2/\text{d}$  iadademstat. At screening, the BM was infiltrated with 40% blasts. After treatment, the neutrophil and platelet counts improved and were normal by D20; the patient exhibited a progressive increase of the absolute neutrophil count (ANC) beginning at D5 (Fig 4A). Induction of *CD86* started on D5, and expression of differentiation genes *VCAN*, *S100A12*, and *LY96* was observed at D12 (Fig 4B-D). The D29 BM demonstrated morphologic remission that was sustained on D52 (both 2% blasts in a hypocellular BM with persistent thrombocytopenia). Of note, on C1D11, the patient developed acute graft-versus-host disease of the liver and was treated with prednisolone



**FIG 1.** Pharmacokinetics (PK) and pharmacodynamics (PD) assessments of iadademstat. (A) Plasma levels of iadademstat were assessed by High-performance liquid chromatography–tandem mass spectrometry (HPLC-MS/MS) in serial samples (day [D]1, D5, D26), trough samples, and washout samples. Mean  $\pm$  SEM plasma levels in cohorts I–VIII and the extension cohort (EC) are shown. (B) PK/PD relationship for *P116* (peptidase inhibitor 16) expression in patients 11 (cohort IV; triangles), 18 (cohort VI; squares), and 22 (cohort VII; dots) during week 1 of treatment (left panel) or washout (right panel). Black arrows indicate dosing occasions. (continued on following page)



and antithymocyte globulin. A single dose of iadademstat was administered on D89 (C2D1). Death occurred in remission on D92 as a result of sepsis for reasons unrelated to iadademstat.

More modest hematologic improvements were also detected in the DE phase in other patients. Patient 9 in cohort III (30  $\mu\text{g}/\text{m}^2/\text{d}$ ) showed a decrease in BM blasts from 38% to 24% after 1 cycle (D29), as did patient 13 in cohort V (60  $\mu\text{g}/\text{m}^2/\text{d}$ ; 51%-36% on C1D29). The modest plasma levels (Data Supplement) were sufficient to induce robust upregulation of differentiation genes *VCAN*, *S100A12*, and *LY96* (Data Supplement).

In the EC, no responses according to International Working Group criteria were observed, but a number of biologically significant changes were seen, including induction of blast cell differentiation and reduction in blast cell burden. Among patients with *MLL* translocations, patient 29 (FAB M4 AML with t[9;11](q21;q23)) showed reductions in BM and PB blast percentages with a concomitant increase in differentiated cells across multiple treatment cycles. The maximum reduction of blasts in the BM was detected on C3D29 (from 95% to 41%) and in PB on C2D29 (from 93% to 44%; Figs 2B and 2C). Induced expression of differentiation genes, including *VCAN*, *S100A12*, *LY96*, *CD86*, and *ITGAM* was detected in blood cells (Fig 3). Although this patient received hydroxycarbamide 500 mg three times daily from C1D6 (as permitted by protocol) and hydroxycarbamide has been reported to potentiate differentiation of other agents,<sup>27</sup> the robust induction of PD biomarkers was already observed on C1D5, in the absence of hydroxycarbamide.

Patients 28 (FAB-M4 AML with t[10;11](q12;q23)) and 36 (FAB-M5 AML with t[11;17](q23;q21)) developed prominent features of morphologic differentiation during treatment. PB smears revealed significant promonocytic and monocytic differentiation, and biomarker analysis showed induction of *CD86*, *ITGAM*, *LY96*, and especially *VCAN* and *S100A12* (Figs 2A and 3). In the former case, the leukocyte count started to increase after 2 weeks of treatment. The patient developed breathlessness on D25 and was treated with antibiotics, intravenous steroids, and hydroxycarbamide. In the latter case, the patient developed fatal respiratory failure

on D5 of treatment, despite prompt treatment with high-dose dexamethasone. Both patients were classified as having drug-induced differentiation syndrome.

There was modest evidence of promonocytic differentiation in patient 31 (FAB-M4 with t[9;11](q21;q23); Fig 2A and 2B), including induction of *LY96* (Fig 3), and in patient 38 (FAB-M2 AML with *MLL*-partial tandem duplication [PTD]), who showed an increase in PB promonocytes from 0% to 10% on C2D23 (Fig 2B). Patient 33 (FAB-M4 AML with t[9;11](q21;q23)) showed a 95% decrease in PB blasts ( $0.2 \times 10^9/\text{L}$  at screening to  $0.01 \times 10^9/\text{L}$  at end of C1). However, the BM blast percentage was stable during treatment, and no morphologic evidence of blast cell differentiation was observed. Differentiation markers upregulated at the end of C1 included *VCAN* and *CD86* (Fig 3).

Blast cells in PB in patient 39 (FAB-M2 AML with *MLL*-PTD) decreased from 75% to 51% on C1D8 (data not shown), although no morphologic differentiation was evident; upregulation of *LY96* and to a minor extent *VCAN* and *CD86* were observed (Fig 3).


Among the four erythroleukemia (FAB M6) patients, patient 35 demonstrated a modest but consistent proportional reduction in BM blasts (Data Supplement). There was no evidence of morphologic differentiation but upregulation of *VCAN* at the end of C1 was observed in blood, and erythrocytic markers were downregulated in BM (Fig 3; Data Supplement). Patient 32 showed stable disease, with a transient blast reduction from 17% to 8% in BM observed between C1D15 and C1D29 of treatment (Figs 2C and 3).

## DISCUSSION

Iadademstat was the first selective inhibitor of LSD1 to enter clinical trials. Our study reveals that iadademstat exhibits approximately linear PK and a half-life of 40-100 hours. Pharmacodynamic analyses demonstrate rapid target engagement. Patients with R/R AML are difficult to treat, with survival times typically in the range of weeks to months. The majority of enrolled patients were > 65 years of age and presented with pancytopenia at screening, making it a particular challenge to discern drug-induced versus disease-related AEs.

**FIG 1.** (Continued). (C) Maximal impact of iadademstat on platelet levels, represented as % inhibition compared with baseline. Individual (symbols) and mean (bar) values are shown for each cohort. Blue symbols represent outlier values that were excluded for calculation of mean. (D) Example of the platelet dynamics in patient 21 (cohort VII), with a predose count of  $149 \times 10^9/\text{L}$ . A time-dependent reduction is followed by rebound. Black bars indicate iadademstat treatment blocks. (E) PK parameters for iadademstat in the EC. Area under the curve ( $\text{AUC}_{0-24\text{h}}$ ) indicates area under the plasma concentration time curve within time 0 to 24 hours (D1: n = 13, D5: n = 13, D26: n = 9).  $C_{\text{max}}$  indicates maximum (peak) plasma drug concentration (D1: n = 14, D5: n = 14, D26: n = 9).  $T_{\text{max}}$  indicates time to reach maximum (peak) plasma concentration (D1: n = 14, D5: n = 14, D26: n = 9).  $\text{AUC}_{\text{inf}}$  indicates area under the plasma concentration time curve from time zero to infinity and half-life (hours; D26: n = 2). Values are shown as mean  $\pm$  standard deviation (median for  $T_{\text{max}}$ ) with range in brackets. CFB, change from baseline; EoT, end of treatment.





**FIG 2.** Morphologic response to treatment with idademstat. (A) Representative images of blood smears showing morphologic differentiation from patient 28 (top) at screening (left) and cycle 1(C1), day 21 (D21) (right) and patient 31 (bottom) at (continued on following page)

Morph. differ. <sup>a</sup>	Blast, % variation <sup>a</sup>	Patient No.	FAB subtype <sup>b</sup>	Time period (hours; range)	Maximum response ( $-\Delta\Delta\text{Cp}$ )												Blasts, % at $t_0$
					VCAN	LYZ	GPR65	S100A12	LY96	CTSG	ANXA2	CRISP9	VIM	CAMSAP2	CD86	ITGAM	
✓	↓ <sup>c</sup>	28 <sup>d</sup>	M4	600-768	6.6	4.9	3.2	7.1	7.0	5.2	3.1	2.6	1.9	4.6	5.3	4.0	52
✓	↓	29	M4	600-768	2.2	0.9	4.8	2.8	5.9	2.3	3.2	4.4	0.7	3.0	3.9	3.7	95
✓	↓	36 <sup>d</sup>	M5a	98-168	9.1	1.2	0.9	5.0	3.3	-3.3	2.6	3.5	0.5	-1.2	2.9	2.3	74
✓	=	31	M4	98-168	1.2	-2.4	-4.4	-3.9	4.1	2.5	-3.2	-2.5	-1.3	2.9	2.8	2.8	81
✗	=	33	M4	600-768	4.3	1.4	6.4	-3.2	-1.1	na	na	na	na	na	6.0	-1.2	7
✗	=	37	M5a	600-768	-2.9	-4.0	-4.8	-3.5	-4.6	-4.7	-1.1	3.3	-2.0	-2.3	-2.8	-1.8	na
✗	n/a	30	M5b	98-168	-1.6	na	na	na	1.6	na	na	na	na	na	na	na	na
✗	↓	39	M2	98-168	2.2	2.3	3.0	-2.4	3.1	1.8	2.0	3.5	2.4	3.7	2.0	1.3	75
✗	↑	41	M2	98-168	2.1	2.2	2.3	-2.1	-2.4	na	1.6	3.1	na	2.5	-0.8	-3.2	3.4
✓	↑	38	M2	600-768	-4.5	-5.5	-4.7	-5.3	-2.6	2.9	-2.8	-5.2	2.8	-3.4	-4.0	-5.4	31
✗	↓ <sup>c</sup>	32	M6a	98-168	-1.2	-1.2	1.5	2.1	1.5	0.7	1.4	1.1	0.6	1.4	-1.3	-1.8	0
✗	↓	35	M6a	600-768	3.7	-3.1	-2.6	-3.2	-2.0	-3.9	2.3	3.9	2.7	2.6	-0.9	-1.6	1
✗	↑	34	M6a	600-768	-2.6	-2.6	-3.4	-3.5	5.1	-3.0	-0.1	5.0	-3.3	2.8	3.8	3.0	na
✗	↑	40	M6a	600-768	2.6	2.5	2.3	-3.6	3.4	-9.1	2.5	-3.3	3.5	4.8	2.8	1.3	20

**FIG 3.** Molecular response to treatment with iadademstat. Relative gene expression levels in nucleated blood cells of a differentiation biomarker panel in the extension cohort (EC). Magenta values show gene upregulation and pink values show gene downregulation. The maximum response and its timing within the treatment period is shown. Data are expressed as  $-\Delta\Delta\text{Cp}$ , calculated relative to expression of the endogenous gene *HPRT1* and to the predose sample. Information on the occurrence of blast cell differentiation in bone marrow (BM) or blood and the percentage variation is also shown. The final column shows blast percentage in peripheral blood at baseline. (a) In bone marrow and/or peripheral blood. (b) Grey background indicates chromosome alterations involving *MLL*; dark grey, *MLL* fusion. (c) Between D5 and D12 of treatment (patient 28) or between D15 and D29 of treatment (patient 32). (d) Differentiation syndrome diagnosed. Morph. differ., morphologic differentiation.

iadademstat was largely well tolerated with a good safety profile. The majority of AEs were as expected for this patient population and included infections and cytopenias, many of which predated the start of treatment. Drug-related AEs such as fatigue, dysgeusia, diarrhea, and anorexia were managed with standard supportive care. Thrombocytopenia, managed with platelet transfusions where necessary, was frequent and an anticipated on-target effect of treatment with an LSD1 inhibitor based on preclinical studies.


Although efficacy was not the main endpoint, there were nonetheless encouraging signs of activity. The most frequent finding was that of induction of differentiation of leukemic blast cells, with responses observed in PD analyses within the first hours or days of treatment. Indeed, gene expression analysis allowed monitoring of the early pharmacologic response to treatment. Induced differentiation was most notable in patients with AML associated with

an *MLL* gene rearrangement: 80% of evaluable patients (4 of 5) exhibited iadademstat-induced morphologic and molecular blast cell differentiation in blood or BM, and the remaining patient with a *MLL* fusion gene exhibited clearance of circulating blasts.

The particular sensitivity of patients with *MLL*-translocated AML to iadademstat may relate to their dependency on the transcription factor GFI1.<sup>16</sup> *GFI1* knockdown in *MLL*-translocated patient blasts robustly induces differentiation<sup>16</sup> and, in addition to inactivating histone demethylase activity, LSD1 inhibitors may also inactivate GFI1 through impeding the physical interaction of LSD1 with GFI1.<sup>16</sup>

In two patients, drug-induced differentiation was vigorous. In patient 28, toward the end of C1, hyperleukocytosis developed with respiratory failure and cellulitis; the syndrome responded to hydroxycarbamide, antibiotics, and

**FIG 2.** (Continued). screening (left) and C1D14 (right; two images from the same slide and patient are shown, separated by a dotted line). Charts show results of morphologic analysis of (B) blood smears and (C) bone marrow smears from selected *MLL*-translocated patients in the extension cohort (EC). HB, hemoglobin; WCC, white cell count; PLT, platelets.



**FIG 4.** Hematologic and biomarker response in patient 16. (A) Absolute neutrophil count (ANC) and platelet dynamics after treatment (cycle 1) and during the washout period in patient 16 of cohort V. Time course of changes in expression of differentiation biomarkers analyzed by quantitative reverse transcriptase–polymerase chain reaction shows that iadademstat induced expression of (B) *CD86*, (C) *VCAN*, (D) *S100A12*, and (E) *LY96* in blood cells of patient 16.

steroids. In the second patient (patient 36), onset of a differentiation syndrome was early (C1D5) and fulminant, resulting in death from respiratory failure despite treatment with high-dose steroid. The severity of the response may be related to the high blast cell count in blood at the start of therapy ( $14.6 \times 10^9/L$ ). More modest features of morphologic or molecular differentiation, or reduction in blast cells, were also observed in 50% of patients (2 of 4) exhibiting an *MLL*-PTD and 50% (2 of 4) of patients with erythroleukemia. Resting periods (2-4 weeks) scheduled between cycles for safety reasons may have allowed progression in some patients (eg, patient 35). Overall, the data

provide clear evidence of the activity of iadademstat as a differentiating agent in patients with AML.

Of particular interest in the DE cohort was the patient in whom iadademstat induced a CRi after a relapse of disease after allogeneic stem cell transplantation. It is unclear whether the remission was induced as a consequence of a leukemia cell intrinsic effect of iadademstat or a noncell intrinsic effect. Pertaining to the latter, inhibitors of LSD1, including iadademstat, induce expression of *CD86*, a protein expressed on antigen-presenting cells that provides costimulatory signals for T-cell activation and survival.<sup>28-31</sup> Expression of *CD86* in murine AML cells stimulated a graft-

versus-leukemia (GVL) effect and survival in a murine allogeneic transplant model.<sup>32</sup> CD86 levels are often low or nonreactive to stimulation in AML.<sup>33</sup> We hypothesize that induction of CD86 in AML blasts by iadademstat might have stimulated a GVL effect raising the possibility that iadademstat may stimulate antileukemic immunity and could be used as an adjunct to immune therapies. LSD1 ablation has recently been reported to enable checkpoint blockade and overcome resistance to anti-programmed death-1 therapy in a mouse melanoma model.<sup>34</sup>

In summary, iadademstat is a well-tolerated compound with a good safety profile without significant extrahematologic toxicity that acts as a potent differentiating agent in AML. Additional LSD1 inhibitors are under early phase evaluation for efficacy in cancer, including INCB059872, bomedemstat (IMG-7289), and CC-90011. In the setting of leukemia, preclinical data suggest that the activity of iadademstat or

other inhibitors of LSD1 may be further enhanced by combinatorial use of all-trans-retinoic acid, azacitidine, rapamycin, BCL2, and DOT1L inhibitors, among others.<sup>23,35-37</sup> Inhibition of LSD1 has also been proposed as an approach to overcome Bromodomain and Extra-Terminal motif (BET) protein inhibitor resistance in AML.<sup>38</sup> In next-phase combination trials, concomitant use of agents with antiproliferative or cytotoxic activity will likely mitigate the risk of differentiation syndrome. Related to that, a phase IIa clinical trial with iadademstat and azacitidine in patients with de novo AML ineligible for intensive chemotherapy and regardless of molecular subtype is ongoing (ALICE study; EudraCT No.: 2018-000482-36) with preliminary data indicating an above-expected proportion of patients achieving CRi and no evidence of differentiation syndrome.<sup>39</sup> Which molecular subtypes of AML are most sensitive to combination antileukemic approaches, including LSD1 inhibitors, remains to be determined.

## AFFILIATIONS

<sup>1</sup>Hospital Vall d'Hebron, Vall D'Hebron Institute of Oncology, Departament de Medicina, Universitat Autònoma de Barcelona, UAB, Barcelona, Spain

<sup>2</sup>Hospital Universitari I Politècnic La Fe, València, Spain

<sup>3</sup>Centro de Investigación Biomédica en Red de Cáncer, Instituto Carlos III, Madrid, Spain

<sup>4</sup>Institut Gustave Roussy, Villejuif Cedex, France

<sup>5</sup>Hospital Universitario Virgen del Rocío, Sevilla, Spain

<sup>6</sup>Instituto de Biomedicina de Sevilla (Instituto de Biomedicina De Sevilla/Consejo Superior De Investigaciones Científicas/Centro de Investigación Biomédica en Red de Cáncer), Universidad de Sevilla, Sevilla, Spain

<sup>7</sup>Centre Hospitalier Universitaire CHU Bordeaux, Hôpital du Haut Lévéque, Pessac, France

<sup>8</sup>Hospitalier Universitaire de Toulouse, Institut Universitaire du Cancer de Toulouse Oncopole, Université Toulouse III Paul Sabatier, Toulouse, France

<sup>9</sup>National Institute for Health Research UCLH Clinical Research Facility, University College London Hospitals NHS Foundation Trust, London, United Kingdom

<sup>10</sup>Oryzon Genomics, Barcelona, Spain

<sup>11</sup>The Christie NHS Foundation Trust, Manchester, United Kingdom

<sup>12</sup>Cancer Research UK Manchester Institute, The University of Manchester, Manchester, United Kingdom

## CORRESPONDING AUTHOR

Tim C. P. Somervaille, MB BS, PhD, Cancer Research UK Manchester Institute, The University of Manchester, Manchester, M20 4GJ, United Kingdom; e-mail: tim.somervaille@cruk.manchester.ac.uk.

## EQUAL CONTRIBUTION

F.B. and T.C.P.S. contributed equally to this work.

## PRIOR PRESENTATION

Presented in part at the 58th Annual Meeting of the American Society of Hematology, December 3-6, 2016, San Diego, CA.

## SUPPORT

Supported by Oryzon Genomics and IPT-2012-0673-010000 of the INNFACTO program of the Spanish Ministry of Economy and Competitiveness, with contribution of Fondo Europeo de Desarrollo Regional (FEDER) from the European Union and CDTI\_CIP-20131005/EUROSTAR\_E18159. T.C.P.S. is supported by Cancer Research UK Grant No. C5759/A20971. R.P. is supported by the National Institute for Health Research, University College London Hospitals, Biomedical Research Centre.

## CLINICAL TRIAL INFORMATION

EudraCT No.: 2013-002447-29

## AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.19.03250>.

## AUTHOR CONTRIBUTIONS

**Conception and design:** Olga Salamero, César Molinero, Tamara Maes, Carlos Buesa, Francesc Bosch, Tim C. P. Somervaille

**Financial support:** Rakesh Popat, Tamara Maes, Carlos Buesa, Tim C. P. Somervaille

**Administrative support:** M. Isabel Arévalo, Tamara Maes

**Provision of study materials or patients:** Olga Salamero, Pau Montesinos, Christophe Willekens, José Antonio Pérez Simón, Arnaud Pigneux, Christian Récher, Rakesh Popat, Cecilia Crespo, Carlos Buesa, Francesc Bosch, Tim C. P. Somervaille

**Collection and assembly of data:** Olga Salamero, Pau Montesinos, Christophe Willekens, José Antonio Pérez-Simón, Arnaud Pigneux, Christian Récher, Rakesh Popat, Cecilia Carpio, César Molinero, Joaquim Vila, Francesc Bosch, Tim C. P. Somervaille

**Data analysis and interpretation:** Olga Salamero, Arnaud Pigneux, Christian Récher, Rakesh Popat, César Molinero, Cristina Mascaró, M. Isabel Arévalo, Tamara Maes, Carlos Buesa, Francesc Bosch, Tim C. P. Somervaille

**Manuscript writing:** All authors

**Final approval of manuscript:** All authors

**Accountable for all aspects of the work:** All authors

## ACKNOWLEDGMENT

The authors thank the patients who participated in this trial, their families, and the co-investigators, nurses, and study coordinators at each of the sites and referral centers. Oryzon Genomics provided iadademstat

and sponsored the study, and worked with investigators to design the study, as well as collect, analyze, and interpret the pharmacokinetics/pharmacodynamics data. The authors also thank Roser Vives Vilagut and the monitors at SynteractHCR.

## REFERENCES

- Vardiman JW, Thiele J, Arber DA, et al: The 2008 revision of the World Health Organization (WHO) classification of myeloid neoplasms and acute leukemia: Rationale and important changes. *Blood* 114:937-951, 2009
- Döhner H, Estey EH, Amadori S, et al: Diagnosis and management of acute myeloid leukemia in adults: Recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood* 115:453-474, 2010
- Döhner H, Weisdorf DJ, Bloomfield CD: Acute myeloid leukemia. *N Engl J Med* 373:1136-1152, 2015
- Nagel G, Weber D, Fromm E, et al: Epidemiological, genetic, and clinical characterization by age of newly diagnosed acute myeloid leukemia based on an academic population-based registry study (AMLSG Bio). *Ann Hematol* 96:1993-2003, 2017
- Burnett AK, Milligan D, Prentice AG, et al: A comparison of low-dose cytarabine and hydroxyurea with or without all-trans retinoic acid for acute myeloid leukemia and high-risk myelodysplastic syndrome in patients not considered fit for intensive treatment. *Cancer* 109:1114-1124, 2007
- Dombret H, Seymour JF, Butrym A, et al: International phase 3 study of azacitidine vs conventional care regimens in older patients with newly diagnosed AML with >30% blasts. *Blood* 126:291-299, 2015
- Welch JS, Petti AA, Miller CA, et al: TP53 and decitabine in acute myeloid leukemia and myelodysplastic syndromes. *N Engl J Med* 375:2023-2036, 2016
- Wouters BJ, Delwel R: Epigenetics and approaches to targeted epigenetic therapy in acute myeloid leukemia. *Blood* 127:42-52, 2016
- Ossenkoppele G, Löwenberg B: How I treat the older patient with acute myeloid leukemia. *Blood* 125:767-774, 2015
- Bhatt VR, Gundabolu K, Koll T, et al: Initial therapy for acute myeloid leukemia in older patients: Principles of care. *Leuk Lymphoma* 59:29-41, 2018
- Toyota M, Kopecky KJ, Toyota MO, et al: Methylation profiling in acute myeloid leukemia. *Blood* 97:2823-2829, 2001
- Di Croce L, Raker VA, Corsaro M, et al: Methyltransferase recruitment and DNA hypermethylation of target promoters by an oncogenic transcription factor. *Science* 295:1079-1082, 2002
- Fathi AT, Abdel-Wahab O: Mutations in epigenetic modifiers in myeloid malignancies and the prospect of novel epigenetic-targeted therapy. *Adv Hematol* 2012: 469592, 2012
- Shen H, Laird PW: Interplay between the cancer genome and epigenome. *Cell* 153:38-55, 2013
- Shi Y, Lan F, Matson C, et al: Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119:941-953, 2004
- Maiques-Diaz A, Spencer GJ, Lynch JT, et al: Enhancer activation by pharmacologic displacement of LSD1 from GF11 induces differentiation in acute myeloid leukemia. *Cell Rep* 22:3641-3659, 2018
- Sprüssel A, Schulte JH, Weber S, et al: Lysine-specific demethylase 1 restricts hematopoietic progenitor proliferation and is essential for terminal differentiation. *Leukemia* 26:2039-2051, 2012
- Lynch JT, Harris WJ, Somervaille TC: LSD1 inhibition: A therapeutic strategy in cancer? *Expert Opin Ther Targets* 16:1239-1249, 2012
- Schenk T, Chen WC, Göllner S, et al: Inhibition of the LSD1 (KDM1A) demethylase reactivates the all-trans-retinoic acid differentiation pathway in acute myeloid leukemia. *Nat Med* 18:605-611, 2012
- Mould DP, McGonagle AE, Wiseman DH, et al: Reversible inhibitors of LSD1 as therapeutic agents in acute myeloid leukemia: Clinical significance and progress to date. *Med Res Rev* 35:586-618, 2015
- Harris WJ, Huang X, Lynch JT, et al: The histone demethylase KDM1A sustains the oncogenic potential of MLL-AF9 leukemia stem cells. *Cancer Cell* 21: 473-487, 2012
- Feng Z, Yao Y, Zhou C, et al: Pharmacological inhibition of LSD1 for the treatment of MLL-rearranged leukemia. *J Hematol Oncol* 9:24, 2016
- Maes T, Mascaró C, Tirapu I, et al: ORY-1001, a potent and selective covalent KDM1A inhibitor, for the treatment of acute leukemia. *Cancer Cell* 33: 495-511.e12, 2018
- Cook N, Hansen AR, Siu LL, et al: Early phase clinical trials to identify optimal dosing and safety. *Mol Oncol* 9:997-1007, 2015
- Wong KM, Capasso A, Eckhardt SG: The changing landscape of phase I trials in oncology. *Nat Rev Clin Oncol* 13:106-117, 2016
- Grimwade D, Hills RK, Moorman AV, et al: Refinement of cytogenetic classification in acute myeloid leukemia: Determination of prognostic significance of rare recurring chromosomal abnormalities among 5876 younger adult patients treated in the United Kingdom Medical Research Council trials. *Blood* 116:354-365, 2010
- Makishima M, Okabe-Kado J, Honma Y: Growth inhibition and differentiation induction in human monoblastic leukaemia cells by 1alpha-hydroxyvitamin D derivatives and their enhancement by combination with hydroxyurea. *Br J Cancer* 77:33-39, 1998
- Lynch JT, Cockerill MJ, Hitchin JR, et al: CD86 expression as a surrogate cellular biomarker for pharmacological inhibition of the histone demethylase lysine-specific demethylase 1. *Anal Biochem* 442:104-106, 2013
- Freeman GJ, Gribben JG, Boussiotis VA, et al: Cloning of B7-2: A CTLA-4 counter-receptor that costimulates human T cell proliferation. *Science* 262:909-911, 1993
- Azuma M, Ito D, Yagita H, et al: B70 antigen is a second ligand for CTLA-4 and CD28. *Nature* 366:76-79, 1993
- Costello R, Cerdan C, Pavon C, et al: The CD2 and CD28 adhesion molecules induce long-term autocrine proliferation of CD4+ T cells. *Eur J Immunol* 23: 608-613, 1993
- Boyer MW, Valleria DA, Taylor PA, et al: The role of B7 costimulation by murine acute myeloid leukemia in the generation and function of a CD8+ T-cell line with potent in vivo graft-versus-leukemia properties. *Blood* 89:3477-3485, 1997
- Whiteway A, Corbett T, Anderson R, et al: Expression of co-stimulatory molecules on acute myeloid leukaemia blasts may effect duration of first remission. *Br J Haematol* 120:442-451, 2003
- Sheng W, LaFleur MW, Nguyen TH, et al: LSD1 ablation stimulates anti-tumor immunity and enables checkpoint blockade. *Cell* 174:549-563.e19, 2018
- Przepolewski A, Wang ES: Inhibitors of LSD1 as a potential therapy for acute myeloid leukemia. *Expert Opin Investig Drugs* 25:771-780, 2016
- Bewersdorf JP, Shallis R, Stahl M, et al: Epigenetic therapy combinations in acute myeloid leukemia: What are the options? *Ther Adv Hematol* 10: 2040620718816698, 2019

37. Deb G, Wingelhofer B, Amaral FM, et al: Pre-clinical activity of combined LSD1 and mTORC1 inhibition in MLL-translocated acute myeloid leukaemia. *Leukemia* 34:1266-1277, 2020
  38. Bell CC, Fennell KA, Chan YC, et al: Targeting enhancer switching overcomes non-genetic drug resistance in acute myeloid leukaemia. *Nat Commun* 10:2723, 2019
  39. Buesa C, Somervaille TC, Arevalo M, et al: ladademstat shows efficacy in elderly AML patients in combination with azacitidine. ALICE trial. *Blood* 134:3839, 2019 (suppl 1)
- 

## ASCO® Career Center

### Search for Oncology Career Opportunities

Visit the ASCO Career Center today to:

- View oncology jobs in your specialty area
- Upload your resume
- Sign up for job alerts
- Get career advice and tips

You can also attend the annual ASCO Career Fair at the ASCO Annual Meeting.

ASCO Career Center is also available via iPhone app. Search jobs at **[careercenter.asco.org](https://careercenter.asco.org)**



**AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST****First-in-Human Phase I Study of Iadademstat (ORY-1001): A First-in-Class Lysine-Specific Histone Demethylase 1A Inhibitor, in Relapsed or Refractory Acute Myeloid Leukemia**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to [www.asco.org/rwc](http://www.asco.org/rwc) or [ascopubs.org/jco/authors/author-center](http://ascopubs.org/jco/authors/author-center).

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](#)).

**Olga Salamero**

**Honoraria:** Celgene, Astellas Pharma, Jazz Pharmaceuticals, Pfizer, Daiichi Sankyo

**Consulting or Advisory Role:** Pfizer, Celgene, Astellas Pharma

**Speakers' Bureau:** Pfizer

**Expert Testimony:** Astellas Pharma, Jazz Pharmaceuticals, Novartis

**Travel, Accommodations, Expenses:** Celgene, Astellas Pharma, Pfizer, Jazz Pharmaceuticals, Novartis

**Pau Montesinos**

**Consulting or Advisory Role:** AbbVie, Pfizer, Shire, Daiichi Sankyo, Novartis, Celgene, Jazz Pharmaceuticals, Roche-Offman

**Speakers' Bureau:** Otsuka, Celgene, Daiichi Sankyo

**Research Funding:** Celgene (Inst), Janssen-Cilag (Inst), Pfizer (Inst)

**Travel, Accommodations, Expenses:** Amgen

**Christophe Willekens**

**Travel, Accommodations, Expenses:** CHUGAI

**José Antonio Pérez-Simón**

**Consulting or Advisory Role:** Novartis, Janssen, Roche, Jazz Pharmaceuticals, Amgen, Novartis, Janssen, Gilead Sciences, Roche, Jazz Pharmaceuticals

**Research Funding:** Novartis, Janssen, Pfizer, Roche, Takeda

**Patents, Royalties, Other Intellectual Property:** Patent licensed by Entourage Bioscience on cannabinoid derivatives

**Travel, Accommodations, Expenses:** Roche, Gilead Sciences, Janssen

**Arnaud Pigneux**

**Consulting or Advisory Role:** Roche, AbbVie, Astellas Pharma, Amgen, Daiichi Sankyo, Jazz Pharmaceuticals, Pfizer

**Travel, Accommodations, Expenses:** Sanofi

**Christian Récher**

**Consulting or Advisory Role:** Celgene, Amgen, Novartis, Jazz Pharmaceuticals, AbbVie, Janssen, Astellas Pharma, MacroGenics, Daiichi Sankyo, Astellas Pharma, Otsuka, Novartis, Incyte, Pfizer, Roche

**Research Funding:** Celgene (Inst), Amgen (Inst), Novartis (Inst), Jazz Pharmaceuticals (Inst), Astellas Pharma (Inst), Chugai Pharma (Inst), Agios (Inst), Daiichi Sankyo (Inst), MaatPharma (Inst), Roche (Inst), AbbVie

**Travel, Accommodations, Expenses:** Incyte, Celgene, Sanofi, Amgen, Novartis, Daiichi Sankyo, Gilead Sciences

**Rakesh Popat**

**Honoraria:** Janssen, Takeda, Celgene, GlaxoSmithKline

**Consulting or Advisory Role:** Takeda, AbbVie, GlaxoSmithKline, Celgene

**Research Funding:** Takeda (Inst)

**Travel, Accommodations, Expenses:** Janssen, Takeda

**Cecilia Carpio**

**Consulting or Advisory Role:** Regeneron

**Speakers' Bureau:** Takeda

**Travel, Accommodations, Expenses:** Gilead Sciences, Novartis

**César Molinero**

**Employment:** Oryzon Genomics

**Leadership:** Oryzon Genomics

**Stock and Other Ownership Interests:** Oryzon Genomics

**Honoraria:** Oryzon Genomics

**Consulting or Advisory Role:** Marketing Y Comunicación Atril

**Uncompensated Relationships:** Oryzon Genomics

**Cristina Mascaró**

**Employment:** Oryzon Genomics

**Joaquim Vila**

**Employment:** Oryzon Genomics

**M. Isabel Arévalo**

**Employment:** Oryzon Genomics

**Tamara Maes**

**Employment:** Oryzon Genomics, Oryzon Genomics (I)

**Leadership:** Oryzon Genomics, Oryzon Genomics (I)

**Stock and Other Ownership Interests:** Oryzon Genomics, Oryzon Genomics (I), Mendelion Lifesciences, Mendelion Lifesciences (I), Palo Biofarma, Palo Biofarma (I)

**Carlos Buesa**

**Employment:** Oryzon Genomics

**Leadership:** Oryzon Genomics (Inst)

**Stock and Other Ownership Interests:** Oryzon Genomics

**Patents, Royalties, Other Intellectual Property:** Inventor in some of Oryzon's patents

**Francesc Bosch**

**Consulting or Advisory Role:** AstraZeneca, Genentech, Janssen-Cilag, Lilly, AbbVie

**Speakers' Bureau:** AbbVie, Janssen, Roche, AstraZeneca

**Tim C. P. Somervaille**

**Honoraria:** Novartis

**Consulting or Advisory Role:** Novartis

**Research Funding:** Imago Biosciences, Cellcentric

No other potential conflicts of interest were reported.