

Table 1. SM. Disease related characteristics, motor and non-motor symptoms, autonomy for activities of daily living and quality of life in PD patients (n=603).

Age	62.7 ± 8.9
Males (%)	59.5
Disease duration (years)	5.7 ± 4.5
L-dopa eq. daily dose (mg)	576.1 ± 421.6
Motor phenotype (%)	
- Tremoric dominant	44.4
- PIGD	40.6
- Indeterminate	15
Hoehn & Yahr	1.9 ± 0.5
- Stage 1 (%)	22.9
- Stage 2 (%)	67.5
- Stage 3 (%)	8.1
- Stages 4 – 5 (%)	1.5
UPDRS-III	22.8 ± 11.2
UPDRS-IV	2.1 ± 2.5
Motor fluctuations (%)	37.1
Dyskinesia (%)	19.7
FOG-Q	3.9 ± 4.6
- Patients with falls (%)	13.8
PD-CRS	90.8 ± 16
- Cognitive impairment (PD-CRS ≤ 84) (%)	32.1
NMSS	46.7 ± 38.2
- Mild NMS burden (NMSS 1-20) (%)	26.9
- Moderate severe burden (NMSS 21-40) (%)	28.8
- Severe NMS burden (NMSS 41-70) (%)	23.2
- Very severe NMS burden (NMSS > 70) (%)	21.1
BDI-II	8.9 ± 7.4
- Depressive symptoms (%)	51.7
- Major depression (%)	16.7
- Minor depression (%)	17.1
- Subthreshold depression (%)	18.2
NPI	6.4 ± 8.3
QUIP-RS	4.5 ± 8.3
PDSS	115 ± 26.3
VAS-PAIN	2.7 ± 2.9
- Patients with pain (%)	56.9
VASF – physical	3.1 ± 2.8
VASF – mental	2.2 ± 2.6
ADLS	88 ± 10.5
- Patients with functional dependency (%)	10.1
PDQ-39SI	17.6 ± 13.7
PQ-10	7.2 ± 1.6
EUROHIS-QOL8	3.8 ± 0.6

The results represent percentages or mean ± SD. Data about H&Y and UPDRS-III are during the OFF state (first thing in the morning without taking medication in the previous 12 hours).

ADLS, Schwab and England Activities of daily living Scale); BDI, Beck Depression Inventory-II; NMSS, Non-Motor Symptoms Scale; NPI, Neuropsychiatric Inventory; PD, Parkinson's disease; PD-CRS, Parkinson's Disease Cognitive Rating Scale; PDSS, Parkinson's Disease Sleep Scale; PIGD, Postural Instability Gait Difficulty; QUIP-RS, Questionnaire for Impulsive-Compulsive Disorders in Parkinson's Disease-Rating Scale; UPDRS, Unified Parkinson's Disease Rating Scale; VAFS, Visual Analog Fatigue Scale; VAS-Pain, Visual Analog Scale-Pain.