

SUPPLEMENTARY MATERIAL

SOURCES OF THE DATA-BASE INCLUDED IN THE MANUSCRIPT

GENERAL REVIEW ARTICLES

Hutchison DC. Alpha 1-antitrypsin deficiency in Europe: geographical distribution of Pi types S and Z. *Respir Med.* 1998;92:367-77.

Blanco I, E Fernández, EF Bustillo. Alpha-1-antitrypsin PI phenotypes S and Z in Europe: an analysis of the published surveys. *Clin Genet* 2001;60:31-41.

Blanco I, EF Bustillo, MC Rodríguez. Distribution of α1-antitrypsin PI S and PI Z frequencies in countries outside Europe: a meta-analysis. *Clin Genet* 2001;60:431-44.

Miravitlles M. Alpha1-anti-trypsin deficiency: epidemiology and incidence. *Respir Med.* 2002;96:205.

de Serres FJ. Worldwide racial and ethnic distribution of alpha1-antitrypsin deficiency: summary of an analysis of published genetic epidemiologic surveys. *Chest*. 2002;122:1818-29.

Luisetti M, Seersholt N. Alpha1-antitrypsin deficiency. 1: epidemiology of alpha1-antitrypsin deficiency. *Thorax*. 2004;59:164-9.

de Serres FJ, Blanco I, Fernández-Bustillo E. Health implications of alpha1-antitrypsin deficiency in Sub-Saharan African countries and their emigrants in Europe and the New World. *Genet Med*. 2005;7:175-84.

Blanco I, de Serres FJ, Fernandez-Bustillo E, Lara B, Miravitlles M. Estimated numbers and prevalence of PI*S and PI*Z alleles of alpha1-antitrypsin deficiency in European countries. *Eur Respir J*. 2006;27:77-84.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI*S and PI*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. *Eur Respir J*. 2006;28:1091-9.

de Serres FJ, Blanco I. Prevalence of α1-antitrypsin deficiency alleles PI*S and PI*Z worldwide and effective screening for each of the five phenotypic classes PI*MS, PI*MZ, PI*SS, PI*SZ, and PI*ZZ: a comprehensive review. *Ther Adv Respir Dis*. 2012;6:277-95.

Blanco I, de Serres FJ, Cárcaba V, Lara B, Fernández-Bustillo E. Alpha-1 Antitrypsin Deficiency PI*Z and PI*S Gene Frequency Distribution Using on Maps of the World by an Inverse Distance Weighting (IDW) Multivariate Interpolation Method. *Hepat Mon.* 2012;12(10 HCC):e7434.

Blanco I, Bueno P, Diego I, Pérez-Holanda S, Casas-Maldonado F, Esquinas C, Miravitles M. Alpha-1 antitrypsin Pi*Z gene frequency and Pi*ZZ genotype numbers worldwide: an update. *Int J Chron Obstruct Pulmon Dis.* 2017;12:561-569.

Blanco I, Bueno P, Diego I, Pérez-Holanda S, Lara B, Casas-Maldonado F, Esquinas C, Miravitles M. Alpha-1 antitrypsin Pi*SZ genotype: estimated prevalence and number of SZ subjects worldwide. *Int J Chron Obstruct Pulmon Dis.* 2017;12:1683-1694.

Blanco I, Diego I, Bueno P, Pérez-Holanda S, Casas-Maldonado F, Miravitles M. Prevalence of α1-antitrypsin PiZZ genotypes in patients with COPD in Europe: a systematic review. *Eur Respir Rev.* 2020;29:200014.

SPECIFIC ARTICLES FOR DIFFERENT COUNTRIES ALPHABETICALLY ORDERED

(Rejected studies appear in italics; selected studies have the first author highlighted in bold)

AFGHANISTAN

Spitsyn VA, Novoradovskii AG, Spitsyna NKh, Parik Iula. [Polymorphism of alpha1-antitrypsin in Pamir populations. Reproductive compensation--possible mechanism for maintaining genetic diversity gor PI genes in human populations]. Genetika 1989;25:2218-24.

Serum protein polymorphisms in four populations of Afghanistan. Rahimi AG, Goedde HW, Flatz G, Kaifie S, Benkmann HG, Delbrück H. Am J Hum Genet. 1977;29:356-60

Goedde HW, Benkmann HG, Flatz G, Bienzle U, Kroeger A. alpha 1-antitrypsin subtypes in the populations of Germany, Ecuador, Afghanistan, Cameroon, and Saudi-Arabia. Zeitschrift fur Morphologie und Anthropologie 1980;70:341-6.

ALBANIA

Scheil HG, Schuh S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. Coll Antropol. 2002;26:403-10.

ANGOLA

Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. Hum Hered 1986;36:238–42.

ARGENTINA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

Sorroche PB, Fernández Acquier M, López Jove O, Giugno E, Pace S, Livellara B, Legal S, Oyhamburu J, Saez MS. Alpha-1 Antitrypsin Deficiency in COPD Patients: A CrossSectional Study. Arch Bronconeumol. 2015;51:539-43.

AUSTRALIA

Ananthakrishnan R, Biegler B, Dennis PM. Alpha 1-antitrypsin phenotypes in paraproteinaemias. Lancet. 1979;1:561-2.

Mulley JC. Distribution of the six alpha 1-antitrypsin (PI) subtypes from a sample of blood donors. Aust J Exp Biol Med Sci 1980: 58: 409–14.

McPhee H, Ananthakrishnan R, Taft LI. Antiproteases and Down's syndrome in an Australian population. J Med Genet 1980: 17: 170–3.

Mulley JC. Distribution of alpha 1-antitrypsin (PI) subtypes from newborns. Aust J Exp Biol Med Sci 1982: 60: 207–9.

Clark P. Alpha-1-protease inhibitor phenotypes in Australia. Hum Hered 1982;32:225–9

Oakeshott JG, Muir A, Clark P et al. Effects of the protease inhibitor (Pi) polymorphism on alpha-1-antitrypsin concentration and elastase inhibitory capacity in human serum. Ann Human Biol 1985: 12: 149–60.

Muir A, Whitfield JB. Alpha-1-antitrypsin (protease inhibitor) phenotypes and longevity. Hum Hered 1990: 40: 22–8.

Huggard PR, West MJ, Summers KM. Alpha 1-antitrypsin deficiency alleles and blood pressure in an Australian population. Clin Exp Pharmacol Physiol 1996: 23: 600–1.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003; 64: 382-97.

AUSTRIA

Böhme A, Cleve H, Schönitzer D, Reissigl H, Kazda S, Müller W. alpha 1-Antitrypsin (Pi) types and subtypes in the Tyrolean population. *Hum Genet*. 1983;63:193-4.

BELGIUM

Kimpen J, Legius E, Bosmans E, et al. Alpha-1-proteinase inhibitor gene frequencies in Belgium. *Gene Geogr* 1990; 4:159–63.

BOLIVIA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

BOSNIA-HERZEGOVINA

Kozarevic D, Laban M, Budimir M, Vojvodic N, Roberts A, Gordon T, McGee DL. Intermediate alpha1-antitrypsin deficiency and chronic obstructive pulmonary disease in Yugoslavia. *Am Rev Respir Dis.* 1978;117:1039-43.

BOTSWANA

Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. Hum Hered 1986;36:238 –42.

BRAZIL

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

Simoes AL, Kompf J, Ritter H, Luckenbach C, Zischler H, Salzano FM. Electrophoretic and isoelectric focusing studies in Brazilian Indians: data on four systems. Hum Biol 1989; 61:427-38.

CAMEROON

Goedde HW, Benkmann HG, Flatz G, Bienzle U, Kroeger A. alpha 1-antitrypsin subtypes in the populations of Germany, Ecuador, Afghanistan, Cameroon, and Saudi-Arabia. Zeitschrift fur Morphologie und Anthropologie 1980;70:341–46.

CANADA

Ashley MJ, Corey P, Chan-Yeung M. Smoking, dust exposure, and serum alpha 1-antitrypsin. Am Rev Respir Dis 1980;121: 783–88.

Chan-Yeung M, Ashley MJ, Corey P et al. Pi phenotypes and the prevalence of chest symptoms and lung function abnormalities in workers employed in dusty industries. Am Rev Respir Dis 1978; 117: 239–45.

Horne SL, Clarke ME, Barnett GD. Alpha-1-antrypsin deficiency: Its effect on pulmonary function in Saskatchewan grain buyers. Occupational Pulmonry Disease: Focus on Grain Dust and Health. New York: Academic Press, 1980: 347–56.

Moroz SP, Cutz E, Cox DW et al. Liver disease associated with alpha1-antitrypsin deficiency in childhood. J Pediatr 1976; 88:19–25.

Talamo RC, Thurlbeck WM. Alpha-antitrypsin Pi types in postmortem blood. Am Rev Respir Dis 1975;112:201–7.

Singh R, Horne SL, Gerrard JW et al. Alpha 1-antitrypsin phenotypes in Canadian Metis. Hum Hered 1983;33:181–3.

Cox DW, Simpson NE, Jantti R. Group-specific component, alpha1-antitrypsin and esterase D in Canadian Eskimos. Hum Hered 1978;28:341–50.

Joly J, Richer G, Boisvert F et al. alpha-1-Antitrypsin phenotypes in French Canadian newborns. Hum Hered 1980;30:1–2.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. Clin Genet 2003;64:382-97.

CAPE VERDE

Spínola C, Brehm A, Spínola H. Alpha-1-antitrypsin deficiency in the Cape Verde islands (Northwest Africa): High prevalence in a sub-Saharan population. Respir Med. 2010;104:1069-72.

CHILE

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

Goedde HW, Rothhammer F, Benkmann HG, Bogdanski P. Ecogenetic studies in Atacameno Indians. Hum Genet 1984;67:343-6.

CHINA

Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. Ann Hum Biol 1990;17:229–34.

Ying QL, Liang ZQ. Allelic frequencies of plasma alpha-1-antitrypsin in Chinese. Sci Sin[B] 1984;27: 161–8.

Ying QL, Zhang ML, Liang CC, et al. Alpha-1-antitrypsin types in five Chinese national minorities. Hum Genet 1985;71:225–6.

Xu JJ, Cui MY, Chen LZ, et al. Polymorphisms of Pi, Hp, ADA and AK in Mongolian, Korean and Zhuang populations of China. Ann Hum Biol 1986;13:245–51.

Lee SS, Lawton JW, Ko KH, Lam KM, Lin CK. alpha-1 antitrypsin phenotypes by isoelectric focusing in a metropolitan southern Chinese population. J Clin Pathol. 2001;54:798-800.

Zhu YJ. Epidemiological survey of chronic obstructive pulmonary disease and alpha-1-deficiency in China. Respirology. 2001;6 Suppl:S13-5.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI*S and PI*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. Eur Respir J. 2006;28:1091-9.

COLOMBIA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

Kirk RL, McDermid EM, Blake NM, Gajdusek DC, Leyshon WC, MacLennan R. Blood group, serum protein and red cell enzyme groups of Amerindian populations in Colombia. Am J Phys Anthropol. 1974;41:301-16.

COSTA RICA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

CUBA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

DEMOCRATIC REPUBLIC OF THE CONGO

Vandeville D, Martin JP, Ropartz C. Alpha 1-antitrypsin polymorphism of a Bantu population: description of a new allele PiL. Humangenetik. 1974;21:33-8.

DENMARK

Thymann M. Distribution of alpha-1-antitrypsin (Pi) phenotypes in Denmark determined by separator isoelectric focusing in agarose gel. Hum Hered 1986;36:19–23.

Dahl M, Nordestgaard BG, Lange P, Vestbo J, Tybjaerg-Hansen A. Molecular diagnosis of intermediate and severe alpha(1)-antitrypsin deficiency: MZ individuals with chronic obstructive pulmonary disease may have lower lung function than MM individuals. Clin Chem 2001;47:56–62.

DOMINICAN REPUBLIC

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

ECUADOR

Goedde HW, Benkmann HG, Flatz G, Bienzle U, Kroeger A. Alpha 1-antitrypsin subtypes in the populations of Germany, Ecuador, Afghanistan, Cameroon, and Saudi-Arabia. Zeitschrift fur Morphologie und Anthropologie 1980;70:341-6.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

EGYPT

Settin A1, El-Bendary M, Abo-Al-Kassem R, El Baz R. Molecular analysis of A1AT (S and Z) and HFE (C282Y and H63D) gene mutations in Egyptian cases with HCV liver cirrhosis. J Gastrointestin Liver Dis. 2006;15:131-5.

Gharib AF, Karam RA, Pasha HF, Radwan MI, Elsawy WH. Polymorphisms of hemochromatosis, and alpha-1 antitrypsin genes in Egyptian HCV patients with and without hepatocellular carcinoma. Gene. 2011;489:98-102.

EL SALVADOR

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

ESTONIA

Uibo R, Parik J, Maaroos H-I, Samloff IM. Alpha-1-antitrypsin allo- and phenotypes in gastric and duodenal ulcer. Scand J Gastroenterol 1991;26:105–8.

Beckman L, Sikström C, Mikelsaar AV, Krumina A, Kucinskas V. a-1-Antitrypsin (Pi) alleles of West European influence in the Baltic Sea region. Hum Hered 1999;49:52–5.

FINLAND

Arnaud P, Koistinen J, Wilson GB, Fudenberg HH. Alpha-1-antitrypsin (Pi) phenotypes in a Finnish population. Scand J Clin Lab Invest 1977;37:339–43.

Häggblom J, Kettunen K, Karjalainen J, Heliövaara M, Jousilahti P, Saarelainen S. Prevalence of PI*Z and PI*S alleles of alpha-1-antitrypsin deficiency in Finland. Eur Clin Respir J. 2015;2:28829.

Fagerhol MK, Eriksson AW, Monn E. Serum Pi types in some Lappish and Finnish populations. Hum Hered 1969;19:360–4.

Fagerhol MK, Gedde-Dahl T Jr. Genetics of the Pi serum types. Family studies of the inherited variants of serum alpha-1-antitrypsin. Hum Hered 1969;19:354–9.

Arvilommi H. Studies on serum Pi- and C3-polymorphism in Finland. Acta Pathol Microbiol Scand 1972; 234:1–50.

FRANCE

Arnaud P, Chapuis-Cellier C, Vittoz P, Creyssel R. Alpha-1-antitrypsin phenotypes in Lyon, France. *Hum Genet* 1977;39:63–8.

Sesboue R, Charlionet R, Vercaigne D, Guimbretiere J, Martin JB. Genetic variants of serum alpha-1-antitrypsin (Pi types) in Bretons. *Hum Hered* 1978;28:280–4.

Morcamp C, Charlionet R, Sesboue R, Lefevre F, Martin JP. Genetic variants of the Pi system in Normans: comparison with some European populations. *Bull Eur Physiopathol Respir* 1980; 16: Suppl., 285–91.

Sesboue R, Martin JP. Alpha-1-antitrypsin (PI) polymorphism in France, with special regard to the PI*Z allele. *Hum Hered* 1991;41:340–6.

Youinou P, Gueguen A, Martin JP, Jouquan J, Salmon D, Ruffie J, Le Menn G, Salmon C. Genetic variants of serum alpha-1-antitrypsin (Pi types) in a Breton population, the Bigoudens. *Hum Hered* 1984;34:114–9.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clinical Genetics* 2003;63:490-509.

Robinet-Levy M, Rieunier M. [Method for the identification of Pi groups: 1st statistics from Languedoc]. Rev Franc Transfus 1972;15:61–72.

GAMBIA

Welch SG, McGregor IA, Williams K. Alpha 1-antitrypsin (pi) phenotypes in a village population from The Gambia, West Africa: Evidence of a new variant occurring at a polymorphic frequency. *Hum Genet* 1980;53:233–5.

GERMANY

Genz T, Martin JP, Cleve H. Classification of alpha1-antitrypsin (Pi) phenotypes by isoelectrofocusing. Distinction of six subtypes of the PiM phenotype. *Hum Genet* 1977; 38:325–32.

Bencze K, Sabatke L, Fruhmann G. Alpha 1-antitrypsin: the PiMM subtypes. Do they play a role in development of chronic obstructive pulmonary diseases? *Chest*. 1980;77:761-3.

Weidinger S, Schwarzfischer F, Cleve H. Pi subtyping by isoelectric focusing: further genetic studies and application to paternity examinations. *Z Rechtsmed* 1980;86:1–7.

Weidinger S, Cleve H, Patutschnick W. Alpha1-antitrypsin: evidence for a fourth PiM allele. Distribution of the PiM subtypes in Southern Germany. *Z Rechtsmed* 1982;88:203–11.

Patzelt D, Schroder H. Genetic-polymorphism of alpha-1-antitrypsin (Pi system) – data on the formal and populationgenetics of the Pi subtypes. *Dtsch Gesundheitsw* 1983;38:119–23.

Schmechta H, Geserick G. [Population genetic studies on alpha-1 antitrypsin polymorphism (Pi-system) in the population of Berlin]. Dtsch Gesundheitsw 1972;27:1089–91.

Thomas K, Hofmann F. [The serum group system Pi. Studies of the methods and frequency in Dresden and vicinity]. Folia Haematol Int Mag Klin Morphol Blutforsch. 1971;95:84-94.

Kellermann G, Walter H. Investigations on the population genetics of the alpha-1-antitrypsin polymorphism. Humangenetik. 1970;10:145-50.

Goedde HW, HG Benkmann. Gene frequencies of red cell adenosine deaminase, adenylate kinase, phosphoglucomutase, acid phosphatase and serum alpha-1 antitrypsin (Pi) in a German population *Humangenetik*, 1970;10:235-43.

GREECE

Fertakis A, Tsourapas A, Douratsos D, Angelopoulos B. Pi phenotypes in Greeks. Hum Hered. 1974;24:313-6.

Scheil HG1, Schuh S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. Coll Antropol. 2002;26:403-10.

GUATEMALA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

GUAYANA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

HAITI

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

HONDURAS

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

HUNGARY

Goedde HW, Benkmann HG, Kriese L, Bogdanski P, Czeizel A, Bères J. Genetic markers among three populations of Hungary. Gene Geogr 1987; 1: 109–20.

ICELAND

Kellermann G, Walter H. Investigations on the population genetics of the alpha-1-antitrypsin polymorphism. Humangenetik. 1970;10:145-50.

INDIA

Kambh MI, Board PG, Kirwood C. Re-evaluation of the proposed interrelationship between thyroxine-binding globulin (TBG) and alpha1-antitrypsin (PI). Clin Chim Acta 1984;139:65–73.

Walter H, Naidu JM, Danke-Hopfe H, et al. Genetic serum protein markers in eight south Indian caste and tribal populations. Z Morphol Anthropol 1993; 79: 355–65.

Reddy PH, Mastana SS. Genetic variation of serum proteins (GC, TF and PI subtypes) in the Baigas of Madhya Pradesh, India. *Anthropol Anz* 1995;53:53–6.

INDONESIA

Constans J, Gouaillard C, Brequet G. Serum protein polymorphism in Bali (Indonesia). *Ann Hum Biol* 1986;13: 537–45.

Kirk RL, McDermid EM, Blake NM, Wight RL, Yap EH, Simons MJ. The distribution of red cell enzyme and serum protein groups in a population of Dani (Pit River, West Irian). Humangenetik 1973; 17: 345–50.

Malcolm LA, Woodfield DG, Blake NM, Kirk RL, McDermid EM. The distribution of blood, serum protein and enzyme groups on Manus Island (Admiralty Islands, New Guinea). Hum Hered 1972; 22: 305–22.

IRAN

Hashemi M, Alavian SM, Ghavami S, de Serres FJ, Salehi M, Doroudi T, Fard AH, Mehrabifar H, Milani B, Shahri SJ. High prevalence of alpha 1 antitrypsin deficient phenotypes in viral hepatitis B infected patients in Iran. *Hepatol Res* 2005;33:292–7.

Walter H, Mohammadzadeh Z, Schuler I, Farhud DD. Serum protein polymorphisms (HP, TF, GC and PI) in four Iranian population samples. *Ann Hum Biol* 1992;19:35–9.

ISRAEL

Nevo S, Kanaaneh H, Cleve H. A genetic study of alpha 1-antitrypsin in an Israeli Arab population, with a new allele: Piv-s. Isr J Med Sci 1982;18:891–3.

Nevo S. Protease inhibitor subtypes in some population groups from Israel. Hum Hered 1987;37:170–81.

Nevo S, Cleve H, Bar-Shani, Joel A., Liron M. Bulgarian Jews in Israel: genetic blood markers. Red-cell antigens, serum proteins and red-cell isozymes. Hum Hered 1989;39:333–44.

Nevo S, Cleve H. PI polymorphism in Israel: report on six Jewish population groups. Am J Med Genet 1991;39:399–403.

Nevo S, Cleve H, Koller A, Eigel E, Patutschnick W, Kanaaneh H, Joel A. Serum protein polymorphisms in Arab Moslems and Druze of Israel: BF, F13B, AHSG, GC, PLG, PI, and TF. Hum Biol 1992;64:587–603.

Ablin JN, Bar-Shira A, Yaron M, Orr-Urtreger A. Candidate-gene approach in fibromyalgia syndrome: association analysis of the genes encoding substance P receptor, dopamine transporter and alpha1-antitrypsin. Clin Exp Rheumatol. 2009;27(5 Suppl 56):S33-8.

ITALY

Piantelli M, Auconi P, Musiani P. Alpha-1-antitrypsin phenotypes in new-borns from Central and Southern Italy. *Hum Hered* 1978;28:468–73.

Massi G, Fabiano A, Ragusa D, Auconi P, Biagini R, Farinato MA, Bandelloni R. Alpha-1-antitrypsin phenotypes and Pi M subtypes in Italy. *Bull Eur Physiopathol Respir* 1980;16, Suppl., 301–6.

Pascali VL, De Mercurio D. Determination of alpha-1-antitrypsin subtypes in the population of Rome: a study in ultrathin-layer isoelectric focusing. *Hum Hered* 1981;31:296–8.

Pittschieder K, Massi G. Alpha-1-antitrypsin deficiency in two population groups in north Italy. *Padiatr Padol* 1988;23:307–11.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clinical Genetics* 2003;63:490-509.

de Serres FJ, Luisetti M, Ferrarotti I, Blanco I, Fernández-Bustillo E. Alpha-1 antitrypsin deficiency in Italy: regional differences of the PIS and PIZ deficiency alleles of alpha-1 antitrypsin deficiency in Italy. *Monaldi Arch Chest Dis* 2005; 63:133-41.

Seefried-Lehmann S-J, Walter H, Danker-Hopfe H, Ricjards O, De Stefano GF. Investigations on the variability of four genetic serum protein markers (HP; TF; CG and PI subtypes) in Italy. Z Morphol Anthropol 1992;79:215–31.

Massi G, Cotumaccio R, Auconi P. Alpha-1-antitrypsin (alpha-1-AT) phenotypes and PiM subtypes in Italy. Evidence of considerable geographic variability. Hum Genet 1982;61:76–7.

Stakisaitis D, Benetis R. Does alpha-1-proteinase inhibitor play a protective role in coronary atherosclerosis? Med Sci Monit 2001;7:701–11.

JAMAICA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

JAPAN

Harada S, Miyake K, Suzuki H, Oda T. New phenotypes of serum alpha1-antitrypsin in Japanese detected by gel slab isoelectric focusing. Hum Genet 1977;38:333–6.

Miyake K, Suzuki H, Oka H, Oda T, Harada S. Distribution of alpha 1-antitrypsin phenotypes in Japanese: description of Pi M subtypes by isoelectric focusing. Jinrui Idengaku Zasshi 1979;24:55–62.

Ohtani H, Saito M. Alpha-1-antitrypsin: frequencies of PiM subtypes and serum concentration in the Japanese population. Hum Hered 1985;35:62–4.

Sebetan IM, Akaishi S. Alpha-1-antitrypsin (PiM) subtypes in Japanese. Forensic Sci Int 1981;18:155–9.

Yuasa I, Suenaga K, Gotoh Y, Ito K, Yokoyama N, Okada K. PI(alpha 1-antitrypsin) polymorphism in the Japanese: confirmation of PI*M4 and description of new PI variants. Hum Genet 1984;67:209–12.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI*S and PI*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. Eur Respir J. 2006;28:1091-9.

JORDAN

Saleh H, Davrinche C, Charlionet R, Rivat C. Alpha-1-antitrypsin phenotypes in a population of Jordan. Hum Hered 1986;36:192–4.

KAZAKHSTAN

Petrishov VN, Lebedeva IA, Shneider Iu V. [Genetic polymorphism of the alpha 1-antitrypsin system in the native population of the Kazakh SSR]. Genetika 1987;23:2257–64.

LATVIA

Beckman L, Sikström C, Mikelsaar AV, Krumina A, Kucinskas V. a-1-Antitrypsin (Pi) alleles of West European influence in the Baltic Sea region. Hum Hered 1999;49:52–55.

LITHUANIA

Beckman L, Siksto m C, Mikelsaar AV, Krumina A, Kucinskas V. a-1-Antitrypsin (PI) alleles of West European influence in the Baltic Sea region. *Hum Hered* 1999;49: 52–5.

Stakisaitis D, Benetis R. Does alpha-1-proteinase inhibitor play a protective role in coronary atherosclerosis? *Med Sci Monit* 2001;7:701–11.

MACEDONIA

Scheil HG, Schuhens S, Huckenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (PI) in the Balkans. *Coll Antropol*. 2002;26:403-10.

Fertakis A, Tsourapas A, Douratsos D, Angelopoulos B. Pi phenotypes in Greeks. Hum Hered. 1974;24:313-6.

MALAYSIA

Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. *Ann Hum Biol* 1990;17:229–34.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimated numbers and prevalence of PI*S and PI*Z deficiency alleles of alpha1-antitrypsin deficiency in Asia. *Eur Respir J*. 2006;28:1091-9.

MALI

Frants RR, Eriksson AW. Reliable classification of six Pi M subtypes by separator isoelectric focusing. Hum Hered 1978;28:201–9.

MEXICO

Pérez-Rubio G, Jiménez-Valverde LO, Ramírez-Venegas A, Camarena Á, Sansores RH, Flores-Trujillo F, Reséndiz-Hernández JM, Falfán-Valencia R. Prevalence of alpha-1 antitrypsin high-risk variants in Mexican mestizo population and their association with lung function values. *Arch Bronconeumol*. 2015;51:80-5.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

MOROCCO

Ezzikouri S, El Feydi AE, El Kihal L, Afifi R, Benazzouz M, Hassar M, Chafik A, Pineau P, Benjelloun S. Prevalence of common HFE and SERPINA1 mutations in patients with hepatocellular carcinoma in a Moroccan population. *Arch Med Res.* 2008;39:236-41.

El Fakir S, Nejjari C, Serhier Z, Tessier JF, Barberger-Gateau P, Dartigues JF, Raherison C. Alpha 1 antitrypsin phenotypes and obstructive airway disease in subjects over 65 years of age: QUID R cohort. Age Ageing. 2011;40:627-9

MOZAMBIQUE

Kellermann G, Walter H. Investigations on the population genetics of the alpha-1-antitrypsin polymorphism. *Humangenetik* 1970;10:145–150.

NAMIBIA

Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. Hum Hered 1986;36:238 –242.

NEPAL

Yuasa I, Saneshige Y, Okamoto N, Ikawa S, Hikita T, Ikebuchi J, Inoue T, Okada K. Distribution of Hp, Tf, Gc and Pi polymorphisms in a Nepalese population. *Hum Hered.* 1983;33:302-6.

NETHERLANDS

Frants RR, Eriksson AW. Reliable classification of six Pi M subtypes by separator isoelectric focusing. *Hum Hered.* 1978;28:201-9.

Hoffman JJML, van den Broek WGM. Distribution of alpha-1-antitrypsin phenotypes in two Dutch population groups. *Hum Genet* 1976;32:43–48.

Klasen EC, Biemond I, Weterman IT. Alpha 1-antitrypsin levels and phenotypes in Crohn's disease in the Netherlands. *Gut* 1980;21:840–842.

Biedmond I, Shelby WS, Jewell DP. Alpha-1-antitrypsin serum concentration and phenotypes in ulcerative colitis. *Digestion* 1984; 29:124–128.

NEW ZEALAND

Janus ED, Joyce PR, Sheat JM, Carrell RW. Alpha-1-antitrypsin variants in New Zealand. *N Z Med J* 1975;82:289–91.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003;64:382-97.

NICARAGUA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

NIGERIA

Olusi SO, Adeyemo AO. Alpha-1-antitrypsin phenotypes, serum immunoglobulins and circulating bacterial antibodies in Nigerians with chronic obstructive pulmonary disease. *East Afr Med J* 1983;60:506-11.

DeCroo S, Kamboh MI, Ferrell RE. Population genetics of alpha-1-antitrypsin polymorphism in US whites, US blacks and African blacks. Hum Hered 1991;41:215–21.

Awotedu AA, Adelaja AB. Alpha-1-antitrypsin and chronic bronchitis in adult Nigerians. Afr J Med Med Sci 1990;19:291–7.

NORWAY

Fagerhol MK, Tenfjord OW. Serum Pi types in some European, American, Asian and African populations. *Acta Pathol Microbiol Scand.* 1968;72:601-8.

Fagerhol MK, Gedde-Dahl T Jr. Serum Pi types in Norwegians. *Acta Pathol Microbial Scand* 1967;70: 421-8.

Moller P, Fagerhol MK. Pi phenotypes in patients with ankylosing spondylitis, acute anterior uveitis, psoriasis and chronic prostatitis. *Clin Genet* 1984; 26: 339–41.

Gulsvik A, Fagerhol MK. Alpha-1-antitrypsin phenotypes and obstructive lung disease in the city of Oslo. *Scand J Respir Dis* 1979; 60: 267–74.

PAKISTAN

Shahid A, Siddiqui AA, Zuberi SJ, Waqar M. Phenotypes of alpha 1 antitrypsin in Karachi, Pakistan. *J Pak Med Assoc* 2000; 50: 374–6.

PANAMA

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

PAPUA NEW GUINEA

Malcolm LA, Woodfield DG, Blake NM, Kirk RL, McDermid EM. The distribution of blood, serum protein and enzyme groups on Manus Island (Admiralty Islands, New Guinea). Hum Hered. 1972;22:305-22.

Kirk RL, McDermid EM, Blake NM, Wight RL, Yap EH, Simons MJ. The distribution of red cell enzyme and serum protein groups in a population of Dani (Pit River, West Irian). Humangenetik. 1973;17:345-50.

PARAGUAY

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

PERU

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

PHILIPPINES

Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. Ann Hum Biol 1990;17:229–34.

POLAND

Titenko-Holland NV, Kowalska A. Alpha-1-antitrypsin (PI) subtypes in Russians and Poles. HumHered 1992;42:384–6.

Walter H, Danker-Hopfe H, Lemmermann M, Lorenz M. Investigation on the variability of four genetic protein markers in Poland. Z Morphol Anthropol 1992;79: 203–14.

Kolwaska A, Rujner J. Polymorphism locus Pi (alpha-1-antitrypsin) of residents in the Poznan Province. Polski Tygodnik Lekarski (Warszawa) 1994;43:195–7.

Kowalska A, Rujner J, Titenko-Holland NV, Pitacik B. Alpha-1-antitrypsin subtypes in Polish newborns. *Hum Hered* 1995;45:351–4.

Chorostowska-Wynimko J, Struniawski R, Popławska B, Borszewska-Kornacka M. The incidence of alpha-1-antitrypsin (A1AT) deficiency alleles in population of Central Poland--preliminary results from newborn screening. *Pneumonol Alergol Pol*. 2012;80:450-3.

Szczeklik AB, Turowska B, Gzerniawska-Mysik B, Opolska B, Nizankowska E. Serum alpha-1 antitrypsin in bronchial asthma. Am Rev Respir Dis 1974;109:487–90.

PORUGAL

Geada HM, Pereira-Albino JA, Manso C. Polimorphism of alpha-1-antitrypsin in a Portuguese population. *Hum Genet* 1976;32:109–13.

Martin JP, Seboue R, Charlionet R, Ropartz C, Pereira MT. Genetic variants of serum alpha-1-antitrypsin (Pi types) in Portuguese. *Hum Hered* 1976;26:310–4.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. *Clinical Genetics* 2003;63:490-509.

Spínola C, Bruges-Armas J, Pereira C, Brehm A, Spínola H. Alpha-1-antitrypsin deficiency in Madeira (Portugal): the highest prevalence in the world. *Respir Med*. 2009;103:1498-502.

Santos-Rosa MA, Robalo-Cordeiro AJA. Alpha-1-proteinase inhibitor phenotypes in Portuguese population. Eur J Respir Dis 1986; 69: Suppl. 146,167–73.

PUERTO RICO

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis*. 2009;71:96-105.

REPUBLIC OF THE CONGO

Pascali VL, Ranalletta D, Spedini G. Antitrypsin and Gc polymorphisms in some populations of Congo: an unusual, highly frequent mutant, PIS, in Bateke and Babenga. *Ann Hum Biol* 1986;13:267–71.

REPUBLIC OF IRELAND

Carroll TP, O'Connor CA, Floyd O, McPartlin J, Kelleher DP, O'Brien G, Dimitrov BD, Morris VB, Taggart CC, McElvaney NG. The prevalence of alpha-1 antitrypsin deficiency in Ireland. *Respir Res.* 2011;12:91.

ROMANIA

Scheil HG, Schuh S, Hockenbeck W, Schmidt HD. Allele frequencies of alpha-1-antitrypsin (Pi) in the Balkans. *Coll Antropol.* 2002;26:403-10.

RUSSIA

Shurkhal AV, Podogas AV, Altukhov YP. Genetic polymorphism and rare variants of alpha-1-antitrypsin in the population of Moscow. *Genetika* 1984;20: 2066-9.

Shurkhal AV, Podogas AV, Dementeva GM, et al. Rare variants of alpha 1 antitrypsin in a group of families with developmental disorders in newborns. *Genetika* 1987;23:1692-8.

Anatonov MP, Petrukhin IS, Voitova TV, Portenko GM. Distribution of Pi phenotypes in normal subjects and subjects with diseases of the respiratory tract. *Lab Delo* 1988;3:23-7.

Kucher AN, Puzyrev VP, Ivanova OF, Xu JJ, Oz RF. Subtypes of serum proteins among Russian inhabitants of the Tomsk region. *Genetika* 1993;29:845-52.

Kravchuk OI, Spitsyn VA, Ginter EK, Makarov SV. Population genetic characteristics of Highland and Meadow Mari: genetic markers. *Hum Genet* 1996;32:570-5.

SAUDI ARABIA

Goedde HW, Benkmann HG, Agarwal DP, Hirth L, Bienzle U, Dietrich M, Hoppe HH, Orlowski J, Kohne E, Kleihauer E. Genetic studies in Saudi Arabia: red cell enzyme, haemoglobin and serum protein polymorphisms. *Am J Phys Anthropol* 1979;50:271-7.

Al-Balla SRS, El-Hazmi MAF, Al-Dalaan AN, Warsy AS. Alpha-1-antitrypsin phenotypes in rheumatoid-arthritis. *Saudi Med J* 1992;13:555-9.

Warsy AS, Elhazmi MAF, Sedrani SH. Alpha-1-antitrypsin phenotypes in Saudi-Arabia - a study in the Central Province. *Ann Saudi Med* 1991;11:159-62.

Warsy AS, el-Hazim MAF, Hamooda H, Kilic N. Alpha-1-antitrypsin: frequencies of Pim subtypes in a Saudi Population. *Saudi Med J* 1991;12:376-9.

SERBIA

Jelic-Ivanovic Z, Spasojevic-Kalimanovska V, Topic A, Spasic S, Petrovic V. Alpha-1-antitrypsin (Pi) polymorphism in Serbia: deviation of Pi M subtype distribution from the Hardy-Weinberg equilibrium. *Gene Geogr* 1994;8:129–35.

SINGAPORE

Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. *Ann Hum Biol* 1990;17:229–34.

SOMALIA

Massi G, Vecchio FM. Alpha-1-antitrypsin phenotypes in a group of newborn infants in Somalia. *Hum Genet* 1977;38:265–9.

SOUTH AFRICA

Dunn DS, Madhoo B, Turnbull R, Jenkins T. Alpha-1-antitrypsin variation in Southern Africa. *Hum Hered* 1986;36:238–42.

Hitzeroth HW, Bissbort S, van den Berg CT, van Niekerk MM, Op't Hof J. Alpha-1-antitrypsin genetic polymorphism in South Africa. *S Afr Med J* 1987;71:755–9.

Walter H, Eberhardt D, Tegeler M, Wiechmann I, Dankerhopfe H, Bhattachaochya SK, Das MK, Mukherjee BN, Saha PN, Das K. HP, GC and PI polymorphisms in sixteen central Indian tribal populations. *Gene Geogr* 1993;7:193–202.

SOUTH KOREA

Lee SS, Lawton JW, Ko KH. Alpha1 antitrypsin phenotypic variability is not associated with ANCA in southern Chinese. *Ann Rheum Dis* 2001;60:725–6.

Saha N. *Alpha 1-protease inhibitor (PI) subtypes in seven populations of East Asia. Ann Hum Biol* 1990;17:229–34.

SPAIN

Vidal R, Miravitles M, Jardí R, Torrella M, Rodríguez-Frías F, Moral P, Vaqué J. [Study of the frequency of different phenotypes of alpha-1-antitrypsin in a population of Barcelona]. *Med Clin (Barcelona)* 1996;107:211–14.

Blanco I, Fernández E, Rodríguez MC, Fernández A. [Allelic frequency of the gene of alpha-1-antitrypsin in the general population in a county in Asturias]. Med Clin (Barcelona) 1999;133:366–70.

Blanco I. A well-designed/conducted study on alpha-1 antitrypsin epidemiology not quoted. Eur Respir J. 2018;51:1702662.

Puertas MJA, Arranz-Peña ML. [Evaluation of phenotypes of alpha-1-antitrypsin in Castilla-Leon: study in a population of newborns]. Rev Clin Esp 1989;184:106–8.

Moral P. [Anthropogenetic study of various haematological polymorphisms on Menorca island]. PhD thesis. University of Barcelona, Barcelona, Spain, 1986.

Blanco I, Fernandez E Alpha1-antitrypsin Pi phenotypes S and Z in Spain: an analysis of the published surveys. Respir Med 2001;95:109-14.

Blanco I, Fernández-Bustillo E, de Serres FJ, Alkassam D, Rodríguez Menéndez C. PI*S and PI*Z alpha 1-antitrypsin deficiency: estimated prevalence and number of deficient subjects in Spain. Med Clin (Barc) 2004; 123:761-5.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in southern Europe: France, Italy, Portugal and Spain. Clinical Genetics 2003;63:490-509.

SURINAME

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. Monaldi Arch Chest Dis. 2009;71:96-105.

SWEDEN

Hjalmarsson K. Distribution of alpha-1-antitrypsin phenotypes in Sweden. Hum Hered 1988;38:27–30.

Sveger T. Liver disease in alpha1-antitrypsin deficiency detected by screening of 200,000 infants. N Engl J Med. 1976;294:1316-21.

SWITZERLAND

Bar W, Kratzer A. Polymorphism of alpha-1-antitrypsin (Pi) in the Swiss population determined by isoelectric focusing with an immobilized pH gradient. *Hum Hered* 1988;38:106–110.

Scheffrahn W, Ziggiotti E. [Electrophoretic alpha 1-antitrypsin variation in the Swiss population]. [Article in German] *Anthropol Anz.* 1982;40:137-43.

TAJIKISTAN

Spitsyn VA, Novoradovskii AG, Spitsyna NKh, Parik Iula. [Polymorphism of alpha1-antitrypsin in Pamir populations. Reproductive compensation--possible mechanism for maintaining genetic diversity gor PI genes in human populations]. Genetika 1989; 25:2218–24.

THAILAND

Pongae P, Schelp FP. Alpha-1-protease inhibitor phenotypes and serum concentrations in Thailand. *Hum Genet* 1980;54: 119–24.

Chongsrisawat V, Jantaradsamee P, Vivatvakin B, Pongpaew P, Poovorawan Y. Alpha 1-antitrypsin phenotype of children with liver diseases in Thailand. Asian Pac J Allergy Immunol 1998;16: 27–30.

Saha N. Alpha 1-protease inhibitor (PI) subtypes in seven populations of east Asia. Ann Hum Biol 1990;17: 229–34.

TRINIDAD AND TOBAGO

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

TUNISIA

Denden S, Zorzetto M, Amri F, Knani J, Ottaviani S, Scabini R, Gorrini M, Ferrarotti I, Campo I, Chibani JB, Khelil AH, Luisetti M. Screening for Alpha 1 antitrypsin deficiency in Tunisian subjects with obstructive lung disease: a feasibility report. *Orphanet J Rare Dis.* 2009;4:12.

Chaabani H, Martin JP, Frants RR, Lefranc G. Genetic study of Tunisian Berbers. II. Alpha 1-antitrypsin (Pi) polymorphism: report of a new allele (Pi S Berber). Exp Clin Immunogenet. 1984;1:19-24.

Chaâbani H, Cox DW. Genetic characterization and origin of Tunisian Berbers. Hum Hered. 1988;38:308-16.

Denden S, Haj Khelil A, Perrin P, Daimi H, Leban N, Ouaja A, Mahdouani K, Hlioui L, Lefranc G, Ben Chibani J. Alpha 1 antitrypsin polymorphism in the Tunisian population with special reference to pulmonary disease. Pathol Biol (Paris). 2008;56:106-10.

UNITED KINGDOM

Cook PJ. The genetics of alpha1-antitrypsin: a family study in England and Scotland. *Ann Hum Genet.* 1975;38:275-87.

Blundell G, Frazer A. Alpha1 antitrypsin phenotypes in Northern Ireland. *Ann Hum Genet* 1975; 38:289–94.

Arnaud P, Galbraith RM, Faulk WP, Black C. Pi phenotypes of alpha1-antitrypsin in Southern England: identification of M subtypes and implications for genetic studies. *Clin Genet* 1979; 15:406–10.

Phillips C, Whitehouse D. A rapid high resolution separator IEF technique for phenotyping alpha-1 antrypsin. *Adv Foren Haemogen* 1990;8: 274–6.

Papiha SS, Pal B, Walker D, Mangion P, Hossain MA. Alpha 1 antitrypsin (PI) phenotypes in two rheumatic diseases: a reappraisal of the association of PI subtypes in rheumatoid arthritis. Ann Rheum Dis. 1989;48:48-52.

Sanders PA, Thomson W, Browne DA, Dyer PA, Grennan DM. alpha 1 Antitrypsin (PI) allotypes in rheumatoid arthritis. Ann Rheum Dis. 1986;45:450-3.

Biemond I, Selby WS, Jewell DP, Klasen EC. Alpha 1-antitrypsin serum concentration and phenotypes in ulcerative colitis. Digestion. 1984;29:124-8.

UNITED STATES

Kueppers F. Serum alpha-1-antitrypsin levels. *N Engl J Med* 1968;279:164–5.

Kueppers F. Alpha1-antitrypsin deficiency. *Birth Defects Orig Artic Ser* 1976;12:213–22.

Kueppers F, Utz G, Simon B. Alpha1-antitrypsin deficiency with M-like phenotype. *J Med Genet* 1977;14:183–6.

Kueppers F, Offord KP. Alpha-1 antitrypsin elevation in healthy neonates. *J Lab Clin Med* 1979;94:475–480.

Roberts A, Kagan A, Rhoads GG et al. Antitrypsin and chronic obstructive pulmonary disease among Japanese-American men. *Chest* 1977;72:489–91.

Lee CC, Kueppers F, Harpel B et al. Alpha-1-antitrypsin (Pi) types in Korean and Chinese populations. *Hum Genet* 1981;57:327–328.

Evans HE, Bognacki NS, Perrott LM et al. Prevalence of alpha 1-antitrypsin Pi types among newborn infants of different ethnic backgrounds. *J Pediatr* 1977;90:621–4.

DeCroo S, Kamboh MI, Ferrell RE. Population genetics of alpha-1-antitrypsin polymorphism in US whites, US blacks and African blacks. *Hum Hered* 1991;41:215–21.

Pierce JA, Eradio B, Dew TA. Antitrypsin phenotypes in St. Louis. *JAMA* 1975; 231:609–12.

Lieberman J, Gaidulis L, Roberts L. Racial distribution of alpha1-antitrypsin variants among junior high school students. *Am Rev Respir Dis* 1976;114:1194–8.

de Serres FJ, Blanco I, Fernández-Bustillo E. Genetic epidemiology of alpha-1 antitrypsin deficiency in North America and Australia/New Zealand: Australia, Canada, New Zealand and the United States of America. *Clin Genet* 2003;64:382-97.

O'Brien ML, Buist NR, Murphrey WH. Neonatal screening for alpha1-antitrypsin deficiency. J Pediatr. 1978;92:1006-10.

Silverman EK, Miletich JP, Pierce JA, Sherman LA, Endicott SK, Broze GJ Jr, Campbell EJ. Alpha-1-antitrypsin deficiency. High prevalence in the St. Louis area determined by direct population screening. Am Rev Respir Dis. 1989;140:961-6.

URUGUAY

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America. *Monaldi Arch Chest Dis.* 2009;71:96-105.

VENEZUELA

Fragoso AT, Barrios GB, Castañeda GC. [The value of the Pi system phenotype in alpha 1-antitrypsin deficiency]. [Spanish]. Acta Gastroenterologica Latinoamericana 1986;16:75-80.

Marini E, Moral P, Petralanda I. Serum protein markers in the Piaroa Indians of Amazonia (Venezuela). Hum Hered 1993;43:232-8.

Fonseca-Perez T, Gonzalez-Coira M, Arias S. PI locus (alpha-1-antitrypsin) allelic frequencies in an Andean Venezuelan population. Gene Geography 1996;10: 65-74.

de Serres FJ, Blanco I, Fernández-Bustillo E. Estimates of PI*S and PI*Z Alpha-1 antitrypsin deficiency alleles prevalence in the Caribbean and North, Central and South America.

Monaldi Arch Chest Dis. 2009;71:96-105.