

ORIGINAL ARTICLE

Phase III, randomized trial of mirvetuximab soravtansine versus chemotherapy in patients with platinum-resistant ovarian cancer: primary analysis of FORWARD I

K. N. Moore^{1*}, A. M. Oza², N. Colombo^{3,4}, A. Oaknin⁵, G. Scambia⁶, D. Lorusso⁷, G. E. Konecny⁸, S. Banerjee⁹, C. G. Murphy¹⁰, J. L. Tanyi¹¹, H. Hirte¹², J. A. Konner¹³, P. C. Lim¹⁴, M. Prasad-Hayes¹⁵, B. J. Monk¹⁶, P. Pautier¹⁷, J. Wang¹⁸, A. Berkenblit¹⁸, I. Vergote¹⁹ & M. J. Birrer²⁰

¹Department of Obstetrics and Gynecology, Stephenson Cancer Center/University of Oklahoma Health Sciences Center, Oklahoma City, USA; ²Division of Medical Oncology and Hematology, Princess Margaret Cancer Centre, Toronto, Canada; ³Department of Gynecologic Oncology, European Institute of Oncology IRCCS; ⁴University of Milan-Bicocca, Milan, Italy; ⁵Medical Oncology Department, Vall d'Hebron University Hospital, Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain; ⁶Gynecology Oncology Unit, Fondazione Policlinico Universitario A. Gemelli IRCCS Roma, Rome; ⁷Gynecologic Oncology, Fondazione IRCCS National Cancer Institute, Milan, Italy; ⁸Department of Obstetrics and Gynecology, University of California Los Angeles, Los Angeles, USA; ⁹Gynaecology, The Royal Marsden NHS Foundation Trust and Institute of Cancer Research, London, UK; ¹⁰Cancer Trials Ireland and Medical Oncology, Bon Secours Hospital, Cork, Ireland; ¹¹Obstetrics and Gynecology, University of Pennsylvania, Philadelphia, USA; ¹²Oncology, Juravinski Cancer Centre, Hamilton, Canada; ¹³Gynecologic Oncology, Memorial Sloan Kettering Cancer Center, New York; ¹⁴Gynecologic Oncology, The Center of Hope Renown Regional Medical Center, Reno; ¹⁵Obstetrics and Gynecology, Icahn School of Medicine at Mount Sinai, New York; ¹⁶Gynecologic Oncology, Arizona Oncology (US Oncology Network), University of Arizona, Creighton University School of Medicine, Phoenix, USA; ¹⁷Medicine, Gustave Roussy, Villejuif, GINECO, Villejuif, France; ¹⁸Clinical Development, ImmunoGen, Inc., Waltham, USA; ¹⁹Department of Obstetrics and Gynecology and Gynecological Oncology, University Hospital Leuven, Leuven Cancer Institute, Leuven, Belgium; ²⁰Comprehensive Cancer Center, University of Alabama at Birmingham, Birmingham, USA



Available online 2 March 2021

Background: Mirvetuximab soravtansine (MIRV) is an antibody-drug conjugate comprising a folate receptor alpha (FR α)-binding antibody, cleavable linker, and the maytansinoid DM4, a potent tubulin-targeting agent. The randomized, open-label, phase III study FORWARD I compared MIRV and investigator's choice chemotherapy in patients with platinum-resistant epithelial ovarian cancer (EOC).

Patients and methods: Eligible patients with 1-3 prior lines of therapy and whose tumors were positive for FR α expression were randomly assigned, in a 2 : 1 ratio, to receive MIRV (6 mg/kg, adjusted ideal body weight) or chemotherapy (paclitaxel, pegylated liposomal doxorubicin, or topotecan). The primary endpoint was progression-free survival [PFS, Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1, blinded independent central review] in the intention-to-treat (ITT) population and in the prespecified FR α high population.

Results: A total of 366 patients were randomized; 243 received MIRV and 109 received chemotherapy. The primary endpoint, PFS, did not reach statistical significance in either the ITT [hazard ratio (HR), 0.98, $P = 0.897$] or the FR α high population (HR, 0.69, $P = 0.049$). Superior outcomes for MIRV over chemotherapy were observed in all secondary endpoints in the FR α high population including improved objective response rate (24% versus 10%), CA-125 responses (53% versus 25%), and patient-reported outcomes (27% versus 13%). Fewer treatment-related grade 3 or higher adverse events (25.1% versus 44.0%), and fewer events leading to dose reduction (19.8% versus 30.3%) and treatment discontinuation (4.5% versus 8.3%) were seen with MIRV compared with chemotherapy.

Conclusions: In patients with platinum-resistant EOC, MIRV did not result in a significant improvement in PFS compared with chemotherapy. Secondary endpoints consistently favored MIRV, particularly in patients with high FR α expression. MIRV showed a differentiated and more manageable safety profile than chemotherapy.

Key words: ovarian cancer, antibody-drug conjugate, folate receptor alpha, mirvetuximab soravtansine, chemotherapy

INTRODUCTION

Epithelial ovarian carcinoma (EOC), including epithelial ovarian, fallopian tube, or primary peritoneal cancer, remains a highly lethal disease, although therapeutic progress has recently been made following the incorporation of molecularly-targeted agents into treatment paradigms.¹ Indeed, the integration of PARP (poly ADP-ribose

*Correspondence to: Dr Kathleen N. Moore, Stephenson Cancer Center at the University of Oklahoma, 800 NE 10th St., Oklahoma City, OK 73104, USA. Tel: +1-405-271-8707
E-mail: kathleen-moore@ouhsc.edu (K.N. Moore).

0923-7534/© 2021 The Author(s). Published by Elsevier Ltd on behalf of European Society for Medical Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

polymerase) inhibitors into the front-line maintenance and recurrent, platinum-sensitive settings,²⁻¹⁰ as well as bevacizumab in all lines of therapy,¹¹⁻¹³ have improved patient outcomes and resulted in an increased prevalence of women living with EOC, despite no real change in incidence or death.¹⁴ Unfortunately, since the vast majority of these patients are not cured with initial therapy, most eventually relapse with disease that is resistant to currently available chemotherapies.¹⁵ Outcomes for patients with platinum-resistant EOC remain particularly poor, with low response rates to further chemotherapy (e.g. 12% in the AURELIA trial),¹³ median progression-free survival (PFS) of 3-4 months,¹⁶ and median overall survival <1 year.¹⁵ Further, subsequent lines of systemic therapy are often associated with cumulative toxicities and limited tolerability for patients. For these reasons, the development of novel therapies for use in the platinum-resistant setting is critical.

One actively pursued molecular target is the cell surface protein, folate receptor alpha (FR α).¹⁷ In contrast to its highly restricted expression in normal tissues, heterogeneous overexpression of FR α is seen in EOC.¹⁸ Its differential distribution pattern, as well as an inherent capacity to internalize large molecules, makes FR α ideally suited for antibody-drug conjugate (ADC)-based therapeutic approaches. ADCs consist of a monoclonal antibody, directed towards tumor-associated antigens, to which a potent cytotoxic agent ('payload') is conjugated via chemical linkage.¹⁹ Moreover, ADCs are a clinically validated class of antineoplastic agents, with nine ADCs currently approved for cancer therapy and >60 others under evaluation in a variety of hematological and solid tumor indications.²⁰

Mirvetuximab soravtansine (MIRV) is an ADC comprising an FR α -binding antibody, cleavable linker, and the maytansinoid DM4, a potent tubulin-targeting agent.²¹ Upon antigen binding, the FR α -ADC complex is rapidly internalized, and DM4 is released. DM4 subsequently elicits potent antimetabolic activity through its ability to suppress microtubule dynamics,²² resulting in cell cycle arrest and apoptosis. Further, the cleavable linker design of MIRV allows active DM4 metabolites to diffuse from antigen-positive tumor cells into neighboring cells and kill them in an antigen-independent manner, an effect known as 'bystander' killing.²³ The phase I clinical experience in patients with platinum-resistant EOC identified the dose, schedule, and target population for a pivotal evaluation of MIRV in this disease setting.²⁴ Here we present results of the phase III FORWARD I trial, designed to assess the safety and clinical activity of MIRV as compared with investigator's choice (IC) chemotherapy in patients with FR α -positive, platinum-resistant EOC.

PATIENTS AND METHODS

Study design and patients

FORWARD I was an open-label, randomized, phase III trial conducted in 12 countries (Trial registration: ClinicalTrials.gov: NCT02631876). The trial was conducted in accordance with the US Food and Drug Administration

regulations, the International Conference on Harmonisation Guidelines for Good Clinical Practice, and the Declaration of Helsinki. The trial was designed by a subset of academic co-authors in collaboration with the sponsor and the Gynecologic Oncology Group Partners (GOG-P), and the protocol was approved by the ethics committee at each participating center.

Eligible patients were 18 years of age or older with histologically confirmed EOC, primary peritoneal cancer, or fallopian tube cancer that was platinum-resistant (defined as progression within 6 months of completion of platinum-containing therapy). Patients were required to have at least one lesion that met the definition of measurable disease by Response Evaluation Criteria in Solid Tumors (RECIST), version 1.1²⁵; have received at least one, but not more than three, prior systemic lines of anticancer therapy. Confirmation of threshold FR α positivity by immunohistochemistry ($\geq 50\%$ of tumor cells with any FR α membrane staining visible at $\leq \times 10$ microscope objective; 50%-74% and $\geq 75\%$ representing medium and high expression, respectively) was determined by central testing using the anti-FOLR1 2.1 antibody at Ventana Medical Systems. Cut-off thresholds for FR α expression level were selected based on the phase I clinical experience with MIRV.²⁴ All patients were required to provide tumor tissue for FR α analysis before enrollment; if archival material was not available, fresh biopsies were allowed using a non-significant risk procedure. Eligible patients also had an Eastern Cooperative Oncology Group performance status score of 0 or 1, and adequate hematologic, renal, and hepatic function. All patients provided written informed consent.

Treatment

Randomization was carried out by means of an interactive online response system with a block design, and stratified according to number of prior therapies, FR α expression, and IC chemotherapy regimen. Patients were randomly assigned, in a 2 : 1 ratio, to receive intravenous infusions of MIRV at a dose of 6 mg/kg (based on adjusted ideal body weight) once every 21 days or IC chemotherapy. All patients treated with MIRV received acetaminophen/paracetamol, dexamethasone, or diphenhydramine 30 min before infusion to prevent infusion-related reactions. Patients were also instructed to self-administer corticosteroid eye drops (1% prednisolone) six times daily on days 1-4 and four times daily on days 5-8 of each cycle during the study and mandated to use preservative-free, lubricating artificial tears on a daily basis (as directed by the product label or treating physician). The chemotherapy regimen selected—paclitaxel, pegylated liposomal doxorubicin (PLD), or topotecan—was stipulated before randomization. Permitted options in the chemotherapy arm were paclitaxel 80 mg/m² on days 1, 8, and 15 of a 4-week treatment cycle, PLD 40 mg/m² on day 1 of a 4-week treatment cycle, or topotecan 4 mg/m² on days 1, 8, and 15 of a 4-week cycle. Alternatively, topotecan could have been administered at 1.25 mg/m² on days 1-5 of a 3-week treatment cycle. The trial

intervention was continued until disease progression on imaging [per RECIST version 1.1, as assessed by blinded independent central review (BICR)], development of unacceptable toxicity, or withdrawal of consent. Crossover was not allowed.

Endpoints and assessments

The primary endpoint was PFS, assessed by BICR, in both the intention-to-treat population (ITT) (all patients who underwent randomization, regardless of the intervention actually received) and in the high FR α subgroup ($\geq 75\%$ of tumor cells with any FR α membrane staining visible at $\leq \times 10$ microscope objective). PFS was defined as the time from randomization to disease progression or time of death from any cause. Tumor response was also assessed by the investigators according to RECIST version 1.1. Computed tomography or magnetic resonance imaging was carried out at baseline and every 6 weeks for the first 36 weeks of study and then every 12 weeks thereafter, until disease progression, discontinuation of study treatment, or death. Secondary endpoints included objective response rate (ORR) by BICR, overall survival (OS), and a patient-reported outcome (PRO) endpoint [defined as number of patients achieving at least a 15% improvement on the QLQ-OV28 abdominal/gastrointestinal symptom subscale (Items 31-36) at week 8/9 assessment]. A hierarchical testing procedure in the order above was used in these key secondary endpoints to control the study-wise type I error. Investigator-assessed PFS and ORR were used as sensitivity analyses to BICR-assessed PFS and ORR. Other secondary endpoints were PFS 2 (PFS2) (defined as the time from randomization to second disease progression or death), duration of response (DOR), and cancer antigen-125 (CA-125) response rate per Gynecologic Cancer Intergroup (GCI) criteria.²⁶

Adverse events were graded with the use of National Cancer Institute Common Terminology Criteria for Adverse Events (CTCAE), version 4.03 and monitored continuously throughout the study from the time of the first dose until 30 days after treatment cessation. Serious adverse events were defined as those that were fatal or life-threatening, required prolonged existing hospitalization, resulted in significant disability or incapacity, or required medical or surgical intervention. Safety data were reviewed by an independent data monitoring committee (IDMC) on a regular basis (approximately every 6 months) both before and after the interim analysis. Ongoing safety data review was carried out by the sponsor and trial investigators.

Statistics

Sample size calculations were based on efficacy assumptions. An interim futility analysis was conducted when 92 PFS events, as assessed by the BICR, had occurred and was reviewed by the IDMC. The observed hazard ratio (HR) was < 1 in all randomized patients as well as in the FR α high expression subgroup, and the study continued as planned. The final analysis was conducted when 244 PFS events were observed. The Hochberg procedure²⁷ was used to control

the study-wise type I error (if the larger of the two P values is < 0.05 then both null hypotheses will be rejected; otherwise, the smaller of the two P values will have to be < 0.025 for the corresponding null hypothesis to be rejected). Assuming a FR α high : medium expression patient ratio of 2 : 1, the study had a power of 85% to 91% to detect a HR of 0.583 in the high FR α subgroup, with the corresponding power for all randomized patients ranging from 75% to 96%. A final OS analysis was conducted 1 year after the final analysis of the primary PFS endpoint. Demographics and baseline characteristics were summarized using descriptive statistics (N , mean, standard deviation, median, and range) for continuous variables and N (%) for discrete variables.

RESULTS

Patients

Between 24 January 2017 and 23 April 2018, 366 patients were randomly assigned to receive MIRV (248) or IC chemotherapy (118). After randomization, 352 patients received at least one dose of their assigned regimen (243 and 109 in the MIRV and IC chemotherapy groups, respectively) and were included in the safety population. The remaining 14 patients did not receive assigned therapy as part of the trial but were included in the ITT population (Figure 1). Baseline characteristics were well balanced between the two treatment groups (Table 1). Most patients had EOC with high-grade serous histology and one or two prior lines of systemic therapy. In total, 218 patients (59.6%) constituted the predefined subset of high FR α expression patients; 147 were assigned to receive MIRV and 71 to receive IC chemotherapy. The median duration of follow-up in the ITT population was 12.5 months for both the MIRV (range, 0.03 to 22.0) and IC chemotherapy groups (range, 0.03 to 20.4). At the time of data cut-off for primary analysis, 27 patients were continuing to receive MIRV (19 of whom were high FR α) and two were still receiving IC chemotherapy (Figure 1).

Efficacy

Analysis of the primary endpoint (PFS, assessed by BICR) was carried out when 244 randomized patients had disease progression or died (Figure 2). For the ITT population, Kaplan–Meier estimates showed no significant difference in PFS between groups [HR, 0.98; 95% confidence interval (CI), 0.73 to 1.31; $P = 0.897$] (Figure 2A); median PFS was 4.1 and 4.4 months for MIRV and IC chemotherapy, respectively. In the prespecified high FR α subgroup (Figure 2B), PFS was longer in patients in the MIRV group compared with IC chemotherapy (median, 4.8 months versus 3.3 months; HR, 0.69, 95% CI, 0.48 to 1.00; $P = 0.049$). However, based on the Hochberg procedure used in the statistical analysis plan for the study, this P value did not meet statistical significance; since the P value for the ITT was > 0.05 , this value was required to be < 0.025 to be significant. Therefore, all P values for primary and secondary




Figure 1. CONSORT diagram.

^a As of 19 February 2019.

PI, principle investigator.

Table 1. Baseline characteristics in the intention-to-treat population		
Characteristic	Mirvetuximab soravtansine (n = 248)	IC chemotherapy (n = 118)
Age, years		
Median	64	64
Range	34-89	31-86
Primary cancer diagnosis		
Epithelial ovarian cancer	207 (83.5)	105 (89.0)
Fallopian tube cancer	14 (5.6)	5 (4.2)
Primary peritoneal cancer	27 (10.9)	8 (6.8)
EOC Histology		
High-grade serous	245 (98.8)	114 (96.6)
Endometrioid	0	1 (0.8)
Serous adenocarcinoma	2 (0.8)	3 (2.5)
Mixed serous and carcinoma	1 (0.4)	0
ECOG PS ^a		
0	141 (56.9)	60 (50.8)
1	106 (42.7)	57 (48.3)
No. of prior systemic therapies ^b		
1 or 2	159 (64.1)	74 (62.7)
3	86 (34.7)	43 (36.4)
FR α expression ^c		
High	147 (59.3)	71 (60.2)
Medium	101 (40.7)	46 (39.0)
Prior exposure		
Paclitaxel	238 (96.0)	113 (95.8)
Bevacizumab	121 (48.8)	55 (46.6)
PARP inhibitor	44 (17.7)	19 (16.1)

Data are number of patients (%) unless indicated otherwise.

ECOG PS, Eastern Cooperative Oncology Group performance status; EOC, epithelial ovarian cancer, IC, investigators choice; PARP, poly ADP-ribose polymerase.

^a Performance status data not available for one patient in each arm.

^b Four patients enrolled were ineligible due to >3 prior lines of therapy.

^c Medium, 50%-74%; High \geq 75% of tumor cells with any FR α membrane staining visible at \leq 10 microscope objective; one patient randomized to the chemotherapy arm was subsequently determined to have a FR α expression level <50%.

endpoints presented hereafter are nominal and no results should be considered as statistically significant.

Secondary endpoint analyses for the ITT (Supplementary Table S1, available at <https://doi.org/10.1016/j.annonc.2021.02.017>) showed the confirmed ORR was higher for MIRV than for IC chemotherapy (22% versus 12%, $P = 0.001$), CA-125 responses were better (51% versus 27%, $P < 0.001$), a longer PFS2 duration was observed (median 10.0 versus 8.4 months; HR, 0.64; 95% CI, 0.49 to 0.84; $P < 0.001$), and there was an improvement in PRO in the EORTC QLQ-OV28 Abdominal/GI Symptom Subscale (32% versus 14%, $P = 0.016$).

Subgroup analyses for survival outcomes are shown in Figure 3. No differences between arms were seen for either PFS or OS in the ITT population (Figures 3A and B); however, the HRs consistently favored MIRV over IC chemotherapy for both these efficacy measures within the high FR α subset (Figures 3C and D). Additional support for superior outcomes for MIRV compared with chemotherapy in this subset of biomarker-defined patients was provided by the secondary endpoint analyses (Supplementary Table S1, available at <https://doi.org/10.1016/j.annonc.2021.02.017>). The confirmed ORR (24% versus 10%, $P = 0.014$), CA-125 response (53% versus 25%, $P = 0.001$), PFS2 (median 10.1 versus 8.4 months; HR, 0.56; 95% CI, 0.39 to 0.79; $P < 0.001$), and PRO improvement (27% versus 13%, $P = 0.143$) were all higher in MIRV-treated group.

No significant difference in OS was seen in the ITT population (Figures 4A and B). In the high FR α subgroup OS was longer with MIRV compared with IC chemotherapy, both at the first cut-off (median, not reached versus 11.8 months;




Figure 2. Kaplan–Meier analysis of progression-free survival (PFS) for the mirvetuximab soravtansine and chemotherapy groups in (A) the intention-to-treat population and (B) in the predefined subset of high FR α patients.

CI, confidence interval.

HR, 0.62; 95% CI, 0.40 to 0.97; $P = 0.033$; Figure 4C) and final analysis (17.3 versus 12.0 months; HR, 0.71; 95% CI, 0.49 to 1.02; $P = 0.063$; Figure 4D), but the differences were not considered statistically significant.

Safety

MIRV was well tolerated, with fewer patients experiencing \geq grade 3 drug-related adverse events (25.1% versus 44.0%), dose reductions (19.8% versus 30.3%), and discontinuations (4.5% versus 8.3%) compared with IC chemotherapy (Table 2). The most common adverse events related to MIRV exposure included nausea (all grades, in 45.7% of patients; grade ≥ 3 in 1%), diarrhea (all grades, in 31.3%; grade ≥ 3 in 2.1%), and fatigue (all grades, in 28.8%; grade ≥ 3 in 1.2%) (Table 2)—with the former two toxicities readily managed

by appropriate supportive measures. Ocular disorders, primarily manifesting as blurred vision (all grades, in 42.0%; grade ≥ 3 in 2.5%) or keratopathy (all grades, 32.5%; grade ≥ 3 in 1.2%), were the most frequent adverse events leading to dose delays and/or reductions (19.8% of patients, Supplementary Table S2, available at <https://doi.org/10.1016/j.annonc.2021.02.017>). In addition to dose modification, these effects were managed by proactive mitigation strategies, and no long-term sequelae were reported. Peripheral neuropathy occurred in 26.7% of patients, with 11.9% of cases being \geq grade 2. The comparative frequency of peripheral neuropathy in the overall chemotherapy population was 18.3%; this was primarily driven by the incidence seen in patients treated with paclitaxel ($n = 32$; all grades, 43.8%, grade ≥ 2 , 28.1%). Alopecia was another adverse event that occurred at a high frequency in




Figure 3. Subgroup analyses.

Progression-free survival (A) and overall survival (B) carried out in the ITT population according to stratification factors and exploratory endpoints at baseline. Progression-free survival (C) and overall survival (D) carried out in predefined subsets of high FR α expression patients according to stratification factors and exploratory endpoints at baseline.

BRCA, breast cancer gene (Foundation Medicine CDx testing); CI, confidence interval; ECOG PS, Eastern Cooperative Oncology Group performance status; FR α , folate receptor alpha; ITT, intention-to-treat; PLD, pegylated liposomal doxorubicin.

paclitaxel-treated patients (21.9%) yet was seen in <1% of patients receiving MIRV. Hematological toxicities observed with MIRV involved cytopenias (all grades, 3.3%-10.7%) that were both lower in prevalence (anemia, 10.7% versus 28.4%; neutropenia, 6.6% versus 39.4%; leukopenia, 3.3% versus 14.7%) as well as severity compared with patients receiving chemotherapy (Table 2).

The percentages of patients experiencing serious adverse events were similar in the MIRV and IC chemotherapy groups (6.6% versus 6.4%; Table 2). Pneumonitis, an adverse event of special interest, occurred in 2.9% of patients (grades 1-3). Two treatment-related deaths due to sepsis occurred in the IC chemotherapy group, one each in patients who received PLD or topotecan. In contrast, no adverse events that emerged during trial intervention, or up to 30 days after the last dose, related to MIRV resulted in death (Table 2).

DISCUSSION

FORWARD I was a randomized, open-label, phase III study comparing MIRV versus IC chemotherapy in patients with platinum-resistant EOC. Baseline characteristics were well balanced between the treatment arms. The study did not meet its primary endpoint of superior PFS in the MIRV arm in the ITT population, nor in the predefined subgroup of FR α high patients.

Based on outcomes observed in corresponding patient populations during the earlier clinical evaluation of MIRV,^{28,29} the negative result was unexpected. In-depth review of the study data revealed that the use of observable membranous staining at $\times 10$ microscope objective ($\times 10$ scoring) as the method of determining FR α positivity for patient enrollment may not have been a reliable method for determining FR α status. In the phase I program for MIRV, the scoring used for eligibility considered not only the proportion score but the intensity of membranous FR α staining as well³⁰ and only samples with sufficient 2+ and 3+ intensity staining (PS2+ scoring) were deemed positive. Further, we have previously reported results from a cohort study evaluating fresh biopsy material that showed FR α expression levels did not significantly shift over time from matched archival samples,³⁰ thus validating the use of archival tissue for determination of eligibility. Exploratory rescoring analyses using PS2+ methodology suggested that use of $\leq \times 10$ scoring allowed enrollment of patients with lower than expected levels of FR α expression, thus diluting the treatment effect of MIRV, in both the ITT and high FR α populations.³¹

Despite this dilution, consistent treatment effects were observed in key secondary endpoints for patients in the protocol-defined high FR α subgroup. In these patients, the confirmed ORR, CA-125 response, PFS2 interval, and PRO improvement all favored MIRV over chemotherapy with




Figure 4. Kaplan–Meier plots of overall survival (OS) in the intention-to-treat population at (A) the time of first data cut-off and (B) final analysis, 1 year after the primary endpoint was reached.

OS estimates for the predefined subset of high FR α patients at first (C) and final analysis (D) are also shown. Hazard ratios (95% CI) are for death and estimated by Cox proportional-hazard regression analysis. CI, confidence interval.

P values below 0.05, with the caveat that these values were not deemed significant due to the statistical analysis plan. Moreover, while FORWARD I was not powered to show a difference in OS, comparatively better OS outcomes were also seen in this subset of patients. Although this was not an unselected population of ovarian cancer patients, the findings additionally add to the body of evidence that elevated FR α expression may be a negative prognostic marker for chemotherapeutic response in EOC.³² FR α high patients responded with limited efficacy to standard chemotherapy (ORR, 10%; median PFS, 3.3 months), further underscoring the critical need for active and tolerable therapies in this biomarker-defined population.

No unexpected toxicities were observed in FORWARD I, and the tolerability profiles of MIRV and chemotherapy were consistent with those from previous studies. MIRV displayed a differentiated safety profile relative to IC chemotherapy, with primarily low-grade nausea, diarrhea, and blurred vision as common adverse events that were readily mitigated with anti-emetics, anti-diarrheals, and lubricating and steroid eye drops. MIRV exposure was associated with significantly less myelosuppression than IC chemotherapy, as evidenced by lower rates of anemia, neutropenia, and thrombocytopenia overall, and no grade 3 or greater neutropenia or thrombocytopenia. Despite its tubulin-directed payload, MIRV was also associated with

less peripheral neuropathy than paclitaxel (12% versus 28% grade ≥ 2) and virtually no alopecia. Patients receiving MIRV required fewer dose reductions or discontinuations due to drug-related adverse events than those on chemotherapy.

Prior efforts to therapeutically target FR α in EOC were hampered by limited single-agent activity, ultimately resulting in negative phase III trials in combinations. Despite negligible activity as monotherapy,³³ the humanized anti-FR α monoclonal antibody farletuzumab showed early promise when administered in combination with chemotherapy, but failed to meet its primary endpoints in randomized phase II or III combination trials in platinum-sensitive EOC^{34,35} and a phase III trial in the platinum-resistant setting was terminated early due to futility.³⁶ A lack of a priori patient selection based on FR α expression level has been suggested to be a contributing factor to the failure of those studies.¹⁷ Vintafolide, a small molecule drug conjugate consisting of a vinca alkaloid linked directly to folate with a short half-life of 20-25 min,³⁷ also displayed limited single-agent activity during its initial clinical evaluation.³⁷ However, this was the first FR α -targeted agent to show a statistically significant improvement, when used in combination, over standard therapy in the randomized phase II PRECEDENT study. This trial evaluated vintafolide in combination with PLD versus PLD alone in women with platinum-resistant ovarian cancer using a companion

Table 2. Summary of treatment-related adverse events in the safety population

Event	Mirvetuximab soravtansine (n = 243)	IC chemotherapy (n = 109)
Any TRAE	230 (94.7)	98 (89.9)
Grade ≥ 3 TRAE	61 (25.1)	48 (44.0)
Serious TRAE	16 (6.6)	7 (6.4)
TRAE leading to dose reduction	48 (19.8)	33 (30.3)
TRAE leading to dose delay	71 (29.2)	31 (28.4)
TRAE leading to discontinuation of trial drug	11 (4.5)	9 (8.3)
TRAE leading to death ^a	0	2 (1.8)

TRAEs ($\geq 15\%$)	Any grade	Grades 3-4	Any grade	Grades 3-4
Any	230 (94.7)	61 (25.1)	98 (89.9)	48 (44.0)
Nausea	111 (45.7)	3 (1.2)	38 (34.9)	0
Vision blurred	102 (42.0)	6 (2.5)	3 (2.8)	0
Keratopathy ^b	79 (32.5)	3 (1.2)	0	0
Diarrhea	76 (31.3)	5 (2.1)	11 (10.1)	0
Fatigue	70 (28.8)	3 (1.2)	34 (31.2)	4 (3.7)
Peripheral neuropathy ^c	65 (26.7)	6 (2.5)	20 (18.3)	3 (2.8)
Dry eye	63 (25.9)	3 (1.2)	2 (1.8)	0
Visual acuity decreased	47 (19.3)	0	1 (0.9)	0
Asthenia	44 (18.1)	2 (0.8)	20 (18.3)	8 (7.3)
Decreased appetite	41 (16.9)	2 (0.8)	9 (8.3)	2 (1.8)
Aspartate aminotransferase increased	40 (16.5)	3 (1.2)	3 (2.8)	1 (0.9)
Vomiting	39 (16.0)	3 (1.2)	16 (14.7)	1 (0.9)
Anemia	26 (10.7)	2 (0.8)	31 (28.4)	12 (11.0)
Constipation	23 (9.5)	0	20 (18.3)	0
Thrombocytopenia	23 (9.5)	0	17 (15.6)	4 (3.7)
Neutropenia	16 (6.6)	0	43 (39.4)	23 (21.1)
Stomatitis	10 (4.1)	0	23 (21.1)	1 (0.9)
Leukopenia	8 (3.3)	0	16 (14.7)	7 (6.4)

Data are number of patients (%).

IC, investigator's choice; PLD, pegylated liposomal doxorubicin; TRAE, treatment-related adverse event.

^a Two deaths observed in the chemotherapy arm involved two cases of grade 5 sepsis seen in patients receiving either PLD or topotecan.

^b Includes preferred terms of keratopathy, keratitis, punctate keratitis, corneal epithelial microcysts, corneal cyst, corneal deposits, limbal stem cell deficiency, corneal disorder, corneal opacity, corneal erosion, corneal pigmentation, keratitis interstitial, and corneal epithelium defect.

^c Includes preferred terms of neuropathy peripheral, peripheral sensory neuropathy, peripheral motor neuropathy, paresthesia, and hypoesthesia.

diagnostic agent known as etarfolatide,³⁸ with the greatest benefit seen in patients whose tumors were 100% positive for FR α expression.³⁹ However, a confirmatory phase III study of the same design was also stopped at interim analysis due to futility. Importantly, MIRV has several potential advantages over these prior modalities, including robust single-agent activity, increased antigen specificity, an extended half-life, and the presence of a cleavable linker which allows for bystander cell killing even in the absence of FR α .¹⁷

The lessons learned from FORWARD I have guided the design of two subsequent studies of MIRV in patients with platinum-resistant, FR α high (by PS2+ scoring) EOC. The single-arm study SORAYA (NCT04296890; prior bevacizumab required) is designed to support accelerated approval, with ORR as the primary endpoint. The randomized MIRASOL trial comparing MIRV monotherapy with IC chemotherapy (NCT04209855) is designed as a

confirmatory study, with PFS as the primary endpoint. MIRASOL incorporates a more efficient and conservative statistical design with 1 : 1 randomization without splitting the alpha and targeting a HR of 0.7.

In conclusion, PFS was not significantly improved with MIRV compared with chemotherapy in FORWARD I. MIRV exhibited a favorable tolerability profile and no new safety signals were observed. The encouraging efficacy outcomes seen in patients with high tumor FR α expression support further work to select patients most likely to benefit from MIRV, as is being done in ongoing clinical trials.

ACKNOWLEDGEMENTS

The authors thank the patients and their families, investigators, co-investigators, and study teams at each of the participating centers.

FUNDING

This work was supported by ImmunoGen, Inc (no grant number).

DISCLOSURE

KNM reports advisory roles for AstraZeneca, AbbVie, Aravive, Eisai, ImmunoGen, GSK/Tesaro, Genentech/Roche, Merck, Mersana, Myriad, Vavotar, and VBL Therapeutics, and research funding from PTC Therapeutics, Lilly, Merck, GSK/Tesaro. DL reports personal fees from Amgen, GSK, PharmaMar, Roche, and Clovis; consulting roles with AstraZeneca, GSK, Merck Sharp & Dohme, Clovis, Amgen, Genmab; and research funding from GSK and Merck. GEK reports personal fees from AstraZeneca, Clovis, and GSK/Tesaro; research funding from Lilly and Merck. HH reports advisory roles and personal fees from AstraZeneca and Merck; research funding from AstraZeneca, GSK, Merck, Roche, and ImmunoGen. BJM reports consulting roles for AbbVie, Amgen, Aravive, AstraZeneca, Clovis, GOG Foundation, Gradalis, ImmunoGen, Laekna Healthcare, Merck, Mersana, Myriad, NuCana, OncoMed, OncoQuest, Pfizer, Roche/Genentech, and Tesaro/GSK. JW and AB report employment with ImmunoGen. All other authors have declared no conflicts of interest.

REFERENCES

- Liu J, Matulonis UA. New strategies in ovarian cancer: translating the molecular complexity of ovarian cancer into treatment advances. *Clin Cancer Res* 2014;20:5150-5156.
- Moore K, Colombo N, Scambia G, et al. Maintenance olaparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med* 2018;379:2495-2505.
- Coleman RL, Oza AM, Lorusso D, et al. Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2017;390:1949-1961.
- Ledermann J, Harter P, Gourley C, et al. Olaparib maintenance therapy in patients with platinum-sensitive relapsed serous ovarian cancer: a preplanned retrospective analysis of outcomes by BRCA status in a randomised phase 2 trial. *Lancet Oncol* 2014;15:852-861.

5. Mirza MR, Monk BJ, Herrstedt J, et al. Niraparib maintenance therapy in platinum-sensitive, recurrent ovarian cancer. *N Engl J Med* 2016;375:2154-2164.
6. Kaufman B, Shapira-Frommer R, Schmutzler RK, et al. Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation. *J Clin Oncol* 2015;33:244-250.
7. Swisher EM, Lin KK, Oza AM, et al. Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial. *Lancet Oncol* 2017;18:75-87.
8. Pujade-Lauraine E, Ledermann JA, Selle F, et al. Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial. *Lancet Oncol* 2017;18:1274-1284.
9. Gonzalez-Martin A, Pothuri B, Vergote I, et al. Niraparib in patients with newly diagnosed advanced ovarian cancer. *N Engl J Med* 2019;381:2391-2402.
10. Ray-Coquard I, Pautier P, Pignata S, et al. Olaparib plus bevacizumab as first-line maintenance in ovarian cancer. *N Engl J Med* 2019;381:2416-2428.
11. Burger RA, Brady MF, Bookman MA, et al. Incorporation of bevacizumab in the primary treatment of ovarian cancer. *N Engl J Med* 2011;365:2473-2483.
12. Coleman RL, Brady MF, Herzog TJ, et al. Bevacizumab and paclitaxel-carboplatin chemotherapy and secondary cytoreduction in recurrent, platinum-sensitive ovarian cancer (NRG Oncology/Gynecologic Oncology Group study GOG-0213): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol* 2017;18:779-791.
13. Pujade-Lauraine E, Hilpert F, Weber B, et al. Bevacizumab combined with chemotherapy for platinum-resistant recurrent ovarian cancer: the AURELIA open-label randomized phase III trial. *J Clin Oncol* 2014;32:1302-1308.
14. National Cancer Institute. Surveillance, Epidemiology, and End Results Program. National Cancer Institute. 2020. Available at: <https://seer.cancer.gov/statfacts/html/ovary.html>. Accessed December 10, 2020.
15. Davis A, Tinker AV, Friedlander M. "Platinum resistant" ovarian cancer: what is it, who to treat and how to measure benefit? *Gynecol Oncol* 2014;133:624-631.
16. Luvero D, Milani A, Ledermann JA. Treatment options in recurrent ovarian cancer: latest evidence and clinical potential. *Ther Adv Med Oncol* 2014;6:229-239.
17. Birrer MJ, Betella I, Martin LP, Moore KN. Is targeting the folate receptor in ovarian cancer coming of age? *Oncologist* 2019;24:425-429.
18. Vergote IB, Marth C, Coleman RL. Role of the folate receptor in ovarian cancer treatment: evidence, mechanism, and clinical implications. *Cancer Metastasis Rev* 2015;34:41-52.
19. Chari RV, Miller ML, Widdison WC. Antibody-drug conjugates: an emerging concept in cancer therapy. *Angew Chem Int Ed Engl* 2014;53:3796-3827.
20. Birrer MJ, Moore KN, Betella I, Bates RC. Antibody-drug conjugate-based therapeutics: state of the science. *J Natl Cancer Inst* 2019;111:538-549.
21. Ab O, Whiteman KR, Bartle LM, et al. IMGN853, a folate receptor-alpha (FRalpha)-targeting antibody-drug conjugate, exhibits potent targeted antitumor activity against FRalpha-expressing tumors. *Mol Cancer Ther* 2015;14:1605-1613.
22. Oroudjev E, Lopus M, Wilson L, et al. Maytansinoid-antibody conjugates induce mitotic arrest by suppressing microtubule dynamic instability. *Mol Cancer Ther* 2010;9:2700-2713.
23. Kovtun YV, Audette CA, Ye Y, et al. Antibody-drug conjugates designed to eradicate tumors with homogeneous and heterogeneous expression of the target antigen. *Cancer Res* 2006;66:3214-3221.
24. Moore KN, Martin LP, O'Malley DM, et al. A review of mirvetuximab soravtansine in the treatment of platinum-resistant ovarian cancer. *Future Oncol* 2018;14:123-136.
25. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228-247.
26. Rustin GJ, Vergote I, Eisenhauer E, et al. Definitions for response and progression in ovarian cancer clinical trials incorporating RECIST 1.1 and CA 125 agreed by the Gynecological Cancer Intergroup (GCIg). *Int J Gynecol Cancer* 2011;21:419-423.
27. Hochberg Y. A sharper Bonferroni procedure for multiple tests of significance. *Biometrika* 1988;75:800-802.
28. Moore KN, Martin LP, O'Malley DM, et al. Safety and activity of mirvetuximab soravtansine (IMGN853), a folate receptor alpha-targeting antibody-drug conjugate, in platinum-resistant ovarian, fallopian tube, or primary peritoneal cancer: a phase I expansion study. *J Clin Oncol* 2017;35:1112-1118.
29. Moore KN, Vergote I, Oaknin A, et al. FORWARD I: a phase III study of mirvetuximab soravtansine versus chemotherapy in platinum-resistant ovarian cancer. *Future Oncol* 2018;14:1669-1678.
30. Martin LP, Konner JA, Moore KN, et al. Characterization of folate receptor alpha (FRalpha) expression in archival tumor and biopsy samples from relapsed epithelial ovarian cancer patients: a phase I expansion study of the FRalpha-targeting antibody-drug conjugate mirvetuximab soravtansine. *Gynecol Oncol* 2017;147:402-407.
31. Moore K, Oza A, Colombo N, et al. FORWARD I (GOG 3011): a phase III study of mirvetuximab soravtansine, a folate receptor alpha (FRa)-targeting antibody-drug conjugate (ADC), versus chemotherapy in patients (pts) with platinum-resistant ovarian cancer (PROC). *Ann Oncol* 2019;30:v403.
32. Toffoli G, Russo A, Gallo A, et al. Expression of folate binding protein as a prognostic factor for response to platinum-containing chemotherapy and survival in human ovarian cancer. *Int J Cancer* 1998;79:121-126.
33. Konner JA, Bell-McGuinn KM, Sabbatini P, et al. Farletuzumab, a humanized monoclonal antibody against folate receptor alpha, in epithelial ovarian cancer: a phase I study. *Clin Cancer Res* 2010;16:5288-5295.
34. Armstrong DK, White AJ, Weil SC, et al. Farletuzumab (a monoclonal antibody against folate receptor alpha) in relapsed platinum-sensitive ovarian cancer. *Gynecol Oncol* 2013;129:452-458.
35. Vergote I, Armstrong D, Scambia G, et al. A randomized, double-blind, placebo-controlled, phase III study to assess efficacy and safety of weekly farletuzumab in combination with carboplatin and taxane in patients with ovarian cancer in first platinum-sensitive relapse. *J Clin Oncol* 2016;34:2271-2278.
36. Sato S, Itamochi H. Profile of farletuzumab and its potential in the treatment of solid tumors. *Onco Targets Ther* 2016;9:1181-1188.
37. Lorusso PM, Edelman MJ, Bever SL, et al. Phase I study of folate conjugate EC145 (vintafolide) in patients with refractory solid tumors. *J Clin Oncol* 2012;30:4011-4016.
38. Maurer AH, Elsinga P, Fanti S, et al. Imaging the folate receptor on cancer cells with 99mTc-etarfolatide: properties, clinical use, and future potential of folate receptor imaging. *J Nucl Med* 2014;55:701-704.
39. Naumann RW, Coleman RL, Burger RA, et al. PRECEDENT: a randomized phase II trial comparing vintafolide (EC145) and pegylated liposomal doxorubicin (PLD) in combination versus PLD alone in patients with platinum-resistant ovarian cancer. *J Clin Oncol* 2013;31:4400-4406.