	Overall				Age			Events	s, n 1	Median, months (95% CI)			
	Overall (n = 152)				•	<75 years (n = 84)						.7 (12.8, 24.9)	
					2	\geq 75 years ($n = 68$)		35		17.0 (9.8, 21.0)		0)	
					•	<80 yea	rs (n = 115)	57		19.1	(15.2, 2	2.1)	
			≥80 years (n = 37)			19		15.3	(8.6, 29	7)			
Kaplan-Meier Estimate	1.0 0.9 0.8 0.7 0.6 0.5 0.4 0.3 0.2 0.1 0.0 0.3 0.3 0.3 0.3	12 15 18 21 Overall surviv		3 36 39 42 45	Kaplan-Meier Estimate	1.0 - 0.9 - 0.8 - 0.7 - 0.6 - 0.5 - 0.4 - 0.3 - 0.2 - 0.1 - 0.0 -	0 3 6	9 12 15 Overal	18 21 24 I survival (n	27 30 nonths)	33 36	39 42	2 45
Dat	ients at risk	0.0.0	ar (months)			ts at ris						10271 10	200
1000000	— 152 137 114 94 7	2 50 33 23	18 12 7 5	5 2 1 1 0	— <75 — ≥75 — <80 — ≥80	years years years years	84 79 66 68 58 48 115104 86 37 33 28	54 42 28 40 30 22 71 57 39 23 15 11	19 15 11 14 8 7 26 20 15 7 3 3	8 6 4 1 5 11 7 1 0	4 2 1 0 5 2 0 0	1	1 0 0 0 1 0 0 0
	Line of therapy		Events, n	Median, months (95% CI)	cher	r platinum-l motherapy t astatic dise	or	Events, n		Mediai (95	n, mon	
	Treatment-naïve		35	17.6 (9.7, 29.7)		Yes	(n = 74)		33		19.8 (1	5.2, 2	4.9)
	1.0 1 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.0 1.	d (n = 83)	41	19.7 (15.0, 21.0)	Kaplan-Meier Estimate	1.0 = 0.9 = 0.8 = 0.7 = 0.6 = 0.5 = 0.4 = 0.3 =	Maryan	-bakenede	سررس	- \			-

Patients at risk

— Yes 74 67 57 51 39 27 18 11 9 6 4 3 2 1 1 0

Figure S6. Overall survival. CI, confidence interval.

Patients at risk

— Treatment-naïve 69 63 51 40 30 21 14 12 9 6 3

— Previously treated 83 74 63 54 42 29 19 11 9 6 4