

Supplementary Table 1. Distribution of symptoms according to overall infected patients and comparison between the 6th SARS-CoV-2 wave and the previous five waves, among Spanish CF children infected between March 2020 and June 2022.

Symptom	Over all n = 351	First 5 waves n = 63	6 th wave n = 288	p-value
Any symptom	264; 75.21%	41; 65,10%	223; 77,4%	.04
Fever	163; 62%	23; 36.5%	140; 48.6%	.08
Headache	54; 20,45%	3; 4.76%	51; 17.7%	.01
Fatigue/Asthenia	44; 16,67%	9; 14.3%	35; 12.15%	.64
Myalgia / Joint pain	38; 14.39%	5; 7.90%	33; 11.45%	.42
Altered cough	139; 52.65%	21; 33.3%	118; 40.9%	.26
Increased sputum	13; 4.92%	8; 12.7%	5; 1.74%	<.001
Dyspnea	10; 3.79%	5; 7.94%	5; 1.74%	.007
Runny nose	63; 23.86%	14; 22.2%	49; 17.0%	.33
Sore throat	30; 11.36%	4; 6.35%	26; 9.03%	.63
Vomiting/Nausea	18; 6.82%	7; 11.1%	11; 3.82%	.017
Diarrhea	13; 4.92%	4; 6.35%	9; 3.12%	.22
Abdominal pain	13; 4.92%	3; 4.76%	10; 3.47%	.62

The proportion of each value was calculated from the non-missing data in each group as depicted by n.