


1984-2014

# Catalan Heart Transplant Registry

## Statistical Report



Generalitat de Catalunya  
**Departament de Salut**



**OCATT**  
Organització Catalana  
de Trasplantaments



# **Catalan Heart Transplant Registry**

Statistical Report 1984-2014

### **Advisory Committee for the Catalan Heart Transplant Registry**

Fèlix Pérez Villa (Hospital Clínic i Provincial de Barcelona)  
Eulàlia Roig (Hospital de la Santa Creu i Sant Pau)  
Nicolás Manito (Hospital Universitari de Bellvitge)  
Dimpna C. Albert (Hospital Universitari Maternoinfantil Vall d'Hebron)  
Nuria Trota (Organització Catalana de Trasplantaments. Servei Català de la Salut)  
Jaume Tort (Organització Catalana de Trasplantaments. Servei Català de la Salut)

### **Notifying Centres**

Hospital Clínic i Provincial de Barcelona  
Hospital de la Santa Creu i Sant Pau  
Hospital Universitari de Bellvitge  
Hospital Universitari Maternoinfantil Vall d'Hebron

**Person responsible for the registry:** Organització Catalana de Trasplantaments

**Registry Management:** Advisory Committee for the Catalan Heart Transplant Registry

**Data collection and entry:** Eulàlia Roig, Sònia Mirabet, Vicens Brossa (Hospital de la Santa Creu i Sant Pau); José Gonzalez Costello, Josep Roca Elies (Hospital Universitari de Bellvitge); Fèlix Pérez Villa, Maria Angeles Castel, Marta Farrero, Ana García (Hospital Clínic i Provincial de Barcelona); Dimpna C. Albert, Ferran Gran (Hospital Universitari Maternoinfantil Vall d'Hebron); Nuria Trota, Pedro López (Organització Catalana de Trasplantaments).

**Data processing and drafting of the report:** Nuria Trota

© Generalitat de Catalunya  
Departament de Salut

Edition: Servei Català de la Salut  
Organització Catalana de Trasplantaments

Barcelona, november 2015

## Acknowledgements

The Catalan Transplant Organization (OCATT), which is responsible for the Catalan Heart Transplant Registry, would like to express its appreciation to all the staff members of the centres authorized to perform heart transplants for their contribution to the maintenance of the registry by supplying data and their participation in the preparation of the report through their contributions.

Dr. Jaume Tort i Bardolet  
Director of OCATT

Address correspondence to:

Nuria Trota  
Catalan Transplant Organization (OCATT)  
Edifici Dr. Frederic Duran i Jordà  
Pg. Taulat 106-116  
08005 Barcelona

Email: [ntrota@catsalut.cat](mailto:ntrota@catsalut.cat)

<http://trasplantaments.gencat.cat>



## Contents

Introduction .....	9
Some Remarks on Methodology.....	10
Evolution of Heart Transplants.....	11
Recipient characteristics .....	13
Donor characteristics .....	20
Transplant characteristics .....	22
Retransplants .....	25
Survival .....	27
Mortality.....	33
Waiting List.....	37
Heart Transplants in Children .....	40





## Introduction

In 1984, the first heart transplant carried out in Catalonia was performed at Hospital de la Santa Creu i Sant Pau. It was also the first successful heart transplant carried out in Spain. A few years later, in 1991, the Hospital Universitari de Bellvitge began working in this field, and was followed by the Hospital Clínic i Provincial de Barcelona in 1998. The Hospital Maternoinfantil Vall d'Hebron was authorized to perform heart and heart-lung transplants in 2006 for children and adolescents.

The Heart Transplant Registry was created in 1993 and contains data on the transplants done in Catalonia since 1984. The data on transplants carried out in the 1984-1993 period were gathered retrospectively, but, since 1994, the registry has systematically gathered data as they have become available.

Publishing the registry is one of the objectives of OCATT, as is managing the data of the Registry Advisory Committee, which responds to the information requirements on planning, resource management and the purchase of services of the Catalan Health Service and the Ministry of Health. The registry is also an information source that is accessible to external users, such as healthcare professionals, and responds to the needs of other sectors. In all cases, processing of and access to data is subject to regulations in force on the protection of personal data.

The main aim of this report is to provide information about the activity and characteristics of the heart transplants carried out in Catalonia in 2014, and to describe the evolution of the transplants carried out since 1984. This information is made available to professionals who are directly involved in this treatment, as well as members of the public administration working in the area of healthcare.

## **Some Remarks on Methodology**

This report describes the evolution of heart transplants in Catalonia and analyses the characteristics of receivers, donors and transplants, as well as the results obtained.

The Kaplan-Meier method was used to calculate the patient survival rate (time before death). The level of statistical significance of the different curves was evaluated using the Log-rank test. The survival curves break off when the number of cases fell below 10.

The probability of receiving a transplantation was calculated bearing in mind the competitive risk model with three events of interest: transplantation, death and removal from the waiting list.

Description of indicators:

### **Annual transplant rate**

The total number of heart transplants carried out during the year at authorized centres, regardless of the place of residence of the receiver, compared with the population of Catalonia (census of 1991, 1996 and, starting in 1997, annual census updates. National Statistics Institute). Expressed per million inhabitants (pmi).

### **Mortality rate at one month**

Percentage of deaths occurring during the 31 days following the heart transplant.

### **Mortality rate at three months**

Percentage of deaths occurring during the 90 days following the heart transplant.

## Evolution of Heart Transplants


In the 1984-2014 period in Catalonia, 1198 heart transplants were performed: 1177 first transplants and 21 retransplants. These transplants were carried out on 1179 patients; two patients received the first transplant outside Catalonia. 1,3% (15) of the transplants were combined with another organ (Table 1).

**Table 1.** Number of combined transplants. 1984-2014

	year activity init	n
Heart – pancreas	1988	1
Heart – kidney	1999	10
Heart – lung	2006	1
Heart – liver	2008	3
<b>Total combined transplants</b>		<b>15</b>

The annual evolution of the number of heart transplants has varied over the years, showing upward trends in 1992 and 1997 (years in which new centres began activity) and until 2000, when the first changes in evolution were registered. In 2014, 56 transplants were performed (Figure 1).


**Figure 1.** Annual evolution of the number of transplants and annual heart transplant rate. 1984-2014



Because of these changes in trend, the annual rate of heart transplants was also affected and showed a clear upward trend in the 1992-2000 period, but has declined since then. In 2014, the transplant rate was 7.4 per million inhabitants (Figure 1).

The annual heart transplant rate varies considerably between countries. Catalonia shows one of the highest rates of transplant activity (Figure 2). However, these data should be interpreted with caution, bearing in mind different factors that affect transplant activity in each country (the healthcare system, indication criteria, population structure, etc.).

**Figure 2.** Heart transplant rate in different countries. 2014



Source: *Newsletter Transplant*. International Figures on Organ Donation and Transplantation - 2014. September 2015. Vol. 20


## Recipient characteristics

### ◆ Sex and age

Of the 1177 patients who received transplants (first transplants) in the 1984-2014 period, 919 (78.1%) were men and 258 (21.9%) were women. In 2014, 41 (74.5%) patients were men and 14 (25.5%) were women.

The mean age of the patients who received their first heart transplant in the 1984-2014 period was 51 (52 for men and 48 for women), the median age was 54 with the range being from age 3 months to 72 years (Figure 3).

**Figure 3.** Number of patients who received their first heart transplant, by age group and sex. 1984-2014



Number of patients (first transplants) = 1177


Over the years, the mean age has increased, going from 48 in 1997 to 53 in 2014 (Figure 4). In 2014, 2 transplants were performed on children under 16 years of age, which have reduced the mean age. Taking into account patients aged 16 or over, the mean age in 2014 was 55.

The increase in mean age over the years is largely due to the gradual increase in transplants done on patients aged 50 or more.

In 2014, 16.4% (9) of the patients who received their first transplantation were between 50 and 60 and 49.1% (27) were over 60. In 1997, these percentages were 33.3% (15) and 20.0% (9), respectively (Figure 5).


Globally, 65.5% of patients were 50 or over when they received their first heart transplant. In fact, 52.3% of all patients were men aged 50 or more.

**Figure 4.** Annual evolution of the mean age of patients receiving their first heart transplant. 1992-2014



Number of patients (first transplants) = 1067

**Figure 5.** Annual evolution of the percentage of patients who received their first heart transplant, by age group. 1992-2014



Number of patients (first transplants) = 1067

#### ◆ Place of residence


91.0% (n = 1073) of patients receiving transplants were residents of Catalonia, 8.6% (n = 101) were residents of another part of Spain, and 0.4% (n = 5) were foreigners. In general, the patients who were not residents of Catalonia came from the Balearic Islands (n = 62) or the autonomous community of Aragon (n = 15).

#### ◆ Indications

The diseases for which a heart transplant is indicated are arranged in four groups: dilated cardiomyopathy, ischemic cardiomyopathy, valvular cardiomyopathy and the “other” category, which includes restrictive cardiomyopathy, congenital cardiomyopathy and hypertrophic cardiomyopathy.


Dilated cardiomyopathy and ischemic cardiomyopathy have been the two most common indications and represent 42.6% and 38.7%, respectively, of all the heart transplants carried out in Catalonia since 1984 (Figure 6). In the case of men, 45.5% of patients suffered from ischemic cardiomyopathy and 39.9% from dilated cardiomyopathy. In the case of women, the most common indication (51.9%) was dilated cardiomyopathy (Figure 7).

**Figure 6.** Percentage of patients who received their first heart transplant, by indication. 1984-2014



Number of patients (first transplants) = 1177

Table 2 provides a breakdown of the indications included in the “Other Indications” category.

**Figure 7.** Percentage of patients who received their first heart transplant, by indication and sex. 1984-2014**Table 2.** Indications included in the “Other Indications” category. 1984-2014


	Men		Women		Global	
	n	%	n	%	n	%
Hypertrophic cardiomyopathy	19	(33,3%)	21	(36,2%)	40	(34,8%)
Restrictive cardiomyopathy	19	(33,3%)	17	(29,3%)	36	(31,3%)
Congenital disease	14	(24,6%)	15	(25,9%)	29	(25,2%)
Arrhythmogenic right ventricular dysplasia	4	(7,0%)	5	(8,6%)	9	(7,8%)
Sarcoidosis	1	(1,8%)	-		1	(0,9%)
<b>Total</b>	<b>57</b>	<b>(100%)</b>	<b>58</b>	<b>(100%)</b>	<b>115</b>	<b>(100%)</b>

In 2014, 43.6% (n = 24) of the patients with ischemic cardiomyopathy, 38.2% (n = 21) with dilated cardiomyopathy and 18.2% (n = 10) with other indications (three patients with restrictive cardiomyopathy, three with hypertrophic cardiomyopathy and four with congenital disease).

The most common disease has changed over the years. Although dilated cardiomyopathy was the most frequent indication in the early years, the percentages of this disease are now much more similar to those of ischemic cardiomyopathy and in some years this indication was even the most frequent. At any rate, given the low number of transplants performed annually, it is difficult to assess the slight changes between one year and another (Figures 8 and 9).




**Figure 8.** Annual evolution of the number of patients who received their first heart transplant, by indication. 1992-2014



Number of patients (first transplants) = 1067


**Figure 9.** Annual evolution of the percentage of patients who received their first heart transplant, by indication. 1992-2014



Number of patients (first transplants) = 1067


In 2004, the registry started using a new system to classify indicated diseases so they could be accounted for more accurately. The most frequent indications in the years from 2004 to 2014 were idiopathic dilated cardiomyopathy, ischemic cardiomyopathy and valvular cardiomyopathy (Figure 10).

**Figure 10.** Percentage of patients who received their first heart transplant, by indication. 2004-2014



The most frequent indication in patients younger than 50 years is dilated cardiomyopathy. In patients over 50 years, the percentage of patients with ischemic cardiomyopathy is slightly higher than dilated cardiomyopathy (Figure 11).

**Figure 11.** Percentage of patients who received their first heart transplant, by indication and age group. 1984-2014



Statistically significant differences in mean age were observed between the four indications ( $p < 0.0001$ ): the patients with ischemic cardiomyopathy or valvular cardiomyopathy were older than those who presented with dilated cardiomyopathy (Table 3). The differences between men and women when treated separately were also statistically significant (Table 4).

**Table 3.** Mean and confidence interval of age, by indication. 1984-2014

	n	mean	IC 95%	range
Dilated cardiomyopathy	501	49	47,5 – 50,2	0 – 72
Ischemic cardiomyopathy	456	55	54,1 – 55,7	1 – 72
Valvular cardiomyopathy	105	55	52,8 – 56,4	17 – 67
Other forms of cardiomyopathy	115	41	37,3 – 44,0	0 – 68
Total	1.177	51	50,1 – 51,7	0 – 72

**Table 4.** Mean and confidence interval of age, by indication and sex. 1984-2014

		n	mean	IC 95%	range
Dilated cardiomyopathy	Men	367	49	47,1 – 50,2	0 – 72
	Women	134	49	46,9 – 52,1	1 – 69
Ischemic cardiomyopathy	Men	418	55	54,3 – 55,9	1 – 72
	Women	38	53	50,1 – 55,7	34 – 69
Valvular cardiomyopathy	Men	77	54	51,7 – 56,1	17 – 67
	Women	28	57	53,2 – 59,9	28 – 65
Other forms of cardiomyopathy	Men	57	43	38,1 – 47,8	1 – 68
	Women	58	38	33,7 – 43,1	0 – 67
Total	Men	919	52	50,8 – 52,5	0 – 72
	Women	258	48	46,3 – 50,2	0 – 69

## Donor characteristics


### ◆ Sex and age

Of the 1085 transplants carried out in the 1992-2014 period, 68.6% (n = 744) of the donors were men and 31.4% (n = 341) were women. In 2014, 62.5% (n = 35) were men and 37.5% (n = 21) were women.

The mean age of the donor over the 1984-2014 period was 34, the median age was also 33 and the range was from some months to 64 years (in the 1992-2014 period, the mean age of the donor was 35 and the median was 35 too). The mean age has increased over the years, going from 25 in 1992 to 45 in 2014 (Figure 12).

This increase has occurred because of older donors, given that in 2014, 51.8% of donors were age 50 and over, whereas there were no donors in this age group in the first years of the programme.

**Figure 12.** Evolution of the mean age of the donor. 1992-2014




Number of donors = 1085

### ◆ Cause of death

The most frequent cause of death of the donor was head trauma (HT), which represented 50.6% (n = 606) of all causes, followed by cerebrovascular accident (CVA) / stroke, which represented 39.5% (n = 473) (In the 1992-2014 period, these percentages were 48.8% and 41.2%, respectively).

In keeping with the increase in the age of donors, the number of donors who died from CVA / stroke also increased (Figure 13). In 2014, 30.4% (n = 17) of donors died from head trauma and 58.9% (n = 33) from CVA / stroke.

**Figure 13.** Evolution of the cause of death of donor (%). 1992-2014




Number of transplants = 1085

◆ **Source of organs**

In the 1984-2014 period, 20.5% (n = 246) of the transplanted organs came from the same hospital where the transplant was carried out, 43.7% (n = 523) from other hospitals in Catalonia, and 35.8% (n = 429) from hospitals outside Catalonia. In 2014, 16.1% (n = 9) of the organs came from the same hospital, 39.3% (n = 22) from Catalonia, and 44.6% (n = 25) from outside Catalonia (Figure 14).

**Figure 14.** Annual evolution of the source of organs (%). 1992-2014



Number of transplants = 1085


## Transplant characteristics

### ◆ Emergency

Of the 1198 transplants carried out in the 1984-2014 period, 23.8% (285) were emergency. The evolution of the percentage of emergency transplants shows an upward trend (Figure 15).

48.2% (27) of the transplants carried out in 2014 were emergency, which is a very similar percentage than in pervious years.


**Figure 15.** Annual evolution of the number and percentage of emergency transplants. 2000-2014



### ◆ Assistance device

In keeping with the increase in emergency transplants, the number of patients who received a ventricular assist device prior to transplant also increased (Figure 16).

In 2014, 77.8% (21) of the emergency transplanted patients received an assist device: 15 intraaortic balloon pump, 2 ECMO, 2 continuous flow device and 2 pulsatile device.

**Figure 16.** Annual evolution of the ventricular assistance device in emergency transplants. 1997-2014

#### ◆ Ischemia time

The mean ischemia time was 178 minutes. Bearing in mind the source of the organ, the differences observed were statistically significant ( $p < 0.0001$ ). When the organ came from a hospital outside Catalonia, the mean ischemia time was 72 minutes longer than when the organ came from a hospital in Catalonia (Table 5).


**Table 5.** Ischemia time, in minutes, in accordance with the source of the organ. 1984-2014

	Same Hospital (n = 245)	Hospital in Catalonia (n = 520)	Hospital outside Catalonia (n = 426)	Total (n = 1191)
Mean	132	157	230	178
Median	127,0	150	232	174
Range	60 – 238	64 – 369	80 – 360	60 – 369
95% CI	128,2 – 136,2	153,5 – 161,2	225,8 – 233,4	174,8 – 181,3

#### ◆ Immunosuppressors

The distribution of immunosuppressors used in the first six weeks after transplant is shown in Figure 17. The overwhelming majority of transplant patients underwent induction immunosuppressive treatment and basiliximab was the drug used most.

**Figure 17.** Immunosuppressant drugs used in the first six weeks after heart transplant. 2004-2014





## Retransplants

A total of 21 of the 1142 heart transplants (1.8%) carried out in the 1984-2014 period have been retransplanted. Two of these patients received the first transplant outside Catalonia.

The time between one transplantation and the other ranged from 1 day to 13 years <sup>1</sup>. The mean was 5 years (the median was 6 years). Specifically, 4 patients (21.1%) received a second transplant within the first week after receiving the first, 3 (15.8%) between the first week and three months after receiving the first transplant, and 12 (63.2%) after the first year.

Tables 6 and 7 show the main characteristics of the retransplants.

**Table 6.** Characteristics of the patients who received a retransplantation, by the time elapsed since the first transplantation. 1984-2014

	0 - 3 months (n = 7)	> 3 months (n = 12)
<b>Sex</b>		
Men	5 (71.4%)	9 (75.0%)
Women	2 (28.6%)	3 (25.0%)
<b>Age (years)</b>		
Mean	44	37
Median	42	39.5
Range	35 – 63	15 – 59
<b>Indications</b>		
Dilated cardiomyopathy	2 (28.6%)	9 (75.0%)
Ischemic cardiomyopathy	3 (42.9%)	3 (25.0%)
Valvular cardiomyopathy	-	-
Other forms of cardiomyopathy	2 (28.6%)	-

Of the 7 patients who received a retransplantation in the first three months after the first transplant, four had died (two by infection) at 31 December 2014. Of the 12 patients who received a retransplantation after the third month, 10 had died at 31 December 2014: 2 due to graft vascular disease, 2 due to primary dysfunction of the graft, 1 due to infection and 5 due to other causes.

<sup>1</sup> Taking into account the patients who received the transplant and retransplant in Catalonia.


**Table 7.** Characteristics of the donor and the transplantation (first transplant), by the time elapsed since the first transplantation. 1984-2014

	0 - 3 months (n = 7)	> 3 months (n = 12)
Donor age (years)		
Mean	20	26
Median	23	22,5
Range	11 – 25	14 – 49
Cause of donor's death		
HT	6 (85,7%)	9 (75,0%)
CVA	-	2 (16,7%)
Other	1 (14,3%)	1 (8,3%)
Ischemia time (minutes)		
Mean	139	168
Median	115	148
Range	82 – 230	95 – 300

## Survival

The survival rate of patients receiving a first heart transplant in Catalonia in the 1984-2014 period was 87% in the first month, 78% in the first year, 72% in the third year, and 69% in the fifth year (Figure 18).

**Figure 18.** Survival rate of patients receiving a heart transplant. 1984-2014



The survival rate percentages were very similar to those of the Spanish and international registries (Table 8).

**Table 8.** Survival: Data comparing the Catalan (RTCC), Spanish (RETC)<sup>2</sup> and international registries (ISHLT)<sup>3</sup>. 1984-2014

	RTCC (1984-2014)	RETC (1984-2014)	ISHLT (1982-6/2013)
1 <sup>st</sup> month	0,86	0,85	0,90
1 <sup>st</sup> year	0,77	0,76	0,82
5 <sup>th</sup> year	0,68	0,65	0,69
10 <sup>th</sup> year	0,55	0,52	0,52
15 <sup>th</sup> year	0,38	0,38	0,35

The overall patient survival rate in the 1984-2014 period was affected by the characteristics of the transplants carried out in the first few years (low number of


<sup>2</sup> González-Vílchez F, et al. Registro Español de Trasplante Cardíaco. XXVI Informe Oficial de la Sección de Insuficiencia Cardíaca y Trasplante Cardíaco de la Sociedad Española de Cardiología (1984-2014). Rev Esp Cardiol. 2015.

<sup>3</sup> International Society for Heart and Lung Transplantation (available at: <http://www.isHLT.org>).

cases, learning period) and by the factors of the transplants carried out in subsequent years (the inclusion of older patients and patients with a more negative prognosis).

The study by period was divided into four time intervals: 1984-1991, 1992-1996, 1997-2001, 2002-2006 and 2007-2014. Statistically significant differences were observed between the five periods ( $p = 0.001$ ), but not between the last four (Figure 19).


**Figure 19.** Survival rate of patients receiving a heart transplant, by period. 1984-2014



In order to gain a more up-to-date look at the results, survival rates were prepared with data on the transplants carried out since 1997. In the 1997-2014 period, the patient survival rate was 89% in the first month, 81% in the first year, 75% in the third year, 71% in the fifth year, and 58% in the tenth year.


The long-term survival rate was slightly lower for men than women (Figure 20), though the differences were not statistically significant ( $p = 0.066$ ).

**Figure 20.** Survival rate of patients receiving a heart transplant, by sex. 1997-2014




Patients over the age of 60 have the lowest survival rate, with statistically significant differences ( $p = 0.030$ ) (Figure 21).

**Figure 21.** Survival rate of patients age 15 and older receiving their first heart transplant, by age group. 1997-2014




Bearing in mind the indicated disease, the patients with valvular cardiomyopathy showed a higher survival rate than patients with ischemic cardiomyopathy, who had the lowest survival rate (Figure 22). The differences between the four diagnostic groups were statistically significant ( $p = 0.015$ ) and also the differences between the two most represented diagnostic categories ( $p = 0.005$ ).

**Figure 22.** Survival rate of patients receiving a heart transplant, by indication. 1997-2014




**Figure 23.** Survival rate of patients receiving a heart transplant, by ischemia time. 1997-2014




When the survival rate was analysed based on the time elapsed between the extraction of the organ and the time of the transplant, it was observed that the probability of survival was greater when the ischemia time was less than or equal to four hours ( $p = 0.004$ ) (Figure 23).

**Figure 24.** Survival rate of patients receiving a heart transplant, by emergency. 1997-2014



**Figure 25.** Survival rate of patients receiving a heart transplant, by emergency and ventricular assistance device. 1997-2014



There are statistically significant differences ( $p = 0.008$ ) in survival rates of patients depending on whether the transplant was an emergency or elective (Figure 24).

Taking into account the different devices used in emergency transplants, the most serious patients have a lower survival rate (Figure 25).




## Mortality

Of the 1179 patients receiving a transplant in the 1984-2014 period, 587 (49.8%) had died at 31 December 2014, 584 (49.5%) remained alive, and monitoring could not be continued on 8 (0.7%).

The most common causes of death were infection (21.6%), primary dysfunction of the graft (14.9%), neoplasm (13.8%) and graft vascular disease (10.4%). The first three causes alone accounted 50% of all deaths (Figure 26).

**Figure 26.** Percentage of deaths, by cause of death. 1984-2014




When the sex of the patient was considered, these four causes of death were the same, though the percentages for each sex varied (Figure 27).

In 27.1% of cases, death occurred in the first month after transplantation (Figure 28) and the main cause of death was primary dysfunction of the graft (Figure 29).

The mortality rate at one month (31 days) is 14.1% (11.9% in 1997-2014 period). In 2014, the mortality rate at one month was 5.4%, lower than the year before (Figure 30).


**Figure 27.** Percentage of deaths, by cause of death and sex. 1984-2014



Number of deaths = 579


In eight cases, the cause of death was unknown

**Figure 28.** Time elapsed between the transplantation and death. 1984-2014




Number of deaths = 587

**Figure 29.** Percentage of deaths by cause of death and time elapsed since the transplantation, patients who received a single heart transplant. 1984-2014




**Figure 30.** Evolution of the mortality rate at first and third months after the heart transplant. 1997-2014



Mortality was analysed in terms of percentage (per 100 people/year) in relation to the period (year of death). The 1984-1991 period was clearly the one with the highest mortality rate (18.06%), whereas in subsequent periods the mortality rate was less than half the rate of the first period (1992-1996: 7.12%; 1997-2001: 6.25%; 2002-2006: 5.45%; 2007-2014: 5.94%). Figure 31 shows the distribution of the cause of

death by period, and reflects the drop in mortality due to primary dysfunction of the graft and severe rejection.

**Figure 31.** Distribution of the cause of death by period, year of death, as a percentage (per 100 people/year). 1984-2014




Number of deaths = 587

In eight cases, the cause of death was unknown

## Waiting List


The number of patients on the waiting list at the end of 2014 was upper than in previous years, going from 27 to 29 (Figure 32).

**Figure 32.** Evolution of the waiting list and the number of heart transplants. 1990-2014



Source: Donor and Transplant Registry

**Figure 33.** Mean number of days on the waiting list to receive a heart transplant. 2000-2014




Source: Donor and Transplant Registry

In 2014, the mean number of days that a patient was on the waiting list for a heart transplant was 150; 202 days in elective transplants and 58 days in emergency transplants (Figure 33).

In 2014, 76 patients were added to the waiting list. Of the patients taken off the list, 6 were removed because their health improved, 2 worsened and one voluntary removal. The mortality rate of the patients on the waiting list was 6.7% (7). Because of the low number of cases in recent years, the major fluctuations observed should be evaluated with caution (Figure 34).

**Figure 34.** Mortality of patients on the waiting list to receive a heart transplant (%). 2000-2014




Source: Donor and Transplant Registry

In the 2000-2014 period, the probability of receiving a heart transplant in the first six months on the waiting list was 62%; in the first year, it was 76% (Figure 35).


When the probability of receiving a transplant was analysed bearing in mind the patient's blood type, it was observed that for patients with blood type A, the probability (62% in the first six months and 74% in the first year) was slightly higher than patients with blood type O (59% in the first six months and 74% in the first year). Patients with blood type B and AB had higher probabilities, but these probabilities were unreliable due to the low number of cases (Figure 36).

**Figure 35.** Probability of receiving a heart transplant. 2000-2014



Source: Donor and Transplant Registry

**Figure 36.** Probability of receiving a heart transplant, by blood type. 2000-2014



Source: Donor and Transplant Registry


## Heart Transplants in Children

Of the 1198 transplants carried out in Catalonia in the 1984-2014 period, 43 were performed on children under 16 years of age (42 first transplants and one retransplant). In 2014, 2 transplants were performed.

Of the 42 patients who received heart transplants, 57.1% (24) were men and 42.9% (18) women. The mean age was 9 (median = 12, range = 3 month – 15 years), although 52.4% (22) of patients were 12-15 years.

The two most frequent indications are dilated cardiomyopathy and congenital disease (Figure 37 and 38).

**Figure 37.** Indications in patients receiving transplants as children (under 16 years of age). 1984-2014




Number of patients (first transplants) = 42

Tables 9 and 10 show the characteristics of the donor and the transplants carried out on children.

Of the 43 patients (under 16 years of age), one needed a retransplant and thirteen had died at 31 December 2014.



**Figure 38.** Indications in patients receiving transplants as children (under 16 years of age) by age group. 1984-2014

Number of patients (first transplants) = 42

**Table 9.** Characteristics of donors in patients receiving transplants as children (under 16 years of age). 1984-2014

<b>Sex of donor</b>	
Men	19 (44,2%)
Women	23 (53,5%)
Missing data	1 (2,3%)
<b>Donor age (years)</b>	
Mean ( $\pm$ SD)	15 ( $\pm$ 12,8)
Median	15
Range	0 – 56
<b>Cause of donor's death</b>	
HT	21 (48,8%)
CVA	5 (11,6%)
Other	17 (39,5%)
<b>Source of organ</b>	
Same hospital	8 (18,6%)
Hospital in Catalonia	7 (16,3%)
Hospital outside Catalonia	28 (65,1%)

**Table 10.** Characteristics of transplants in patients receiving transplants as children (under 16 years of age). 1984-2014

Emergency	
Urgent	21 (48,8%)
Elective	22 (51,2%)
Ischemia time (minutes)	
Mean ( $\pm$ SD)	205 ( $\pm$ 57,7)
Median	206
Range	60 – 340



