

TableS1. Participants' profile

| | TOTAL (n=148) | France (n=37) | Germany (n=10) | Italy (n=31) | Spain (n=37) | UK (n=33) | p value |
|--|--------------------------|--------------------------|---------------------------|-------------------------|-------------------------|----------------------|----------------|
| Medical specialty: neurology | 148 (100%) | 37 (100%) | 10 (100%) | 31 (100%) | 37 (100%) | 33 (100%) | >0.999 |
| Experience in treating migraine | | | | | | | <0.001 |
| 2-5 years | 23 (15.5%) | 3 (8.1%) | - | 8 (25.8%) | 12 (32.5%) | - | |
| 5-10 years | 24 (16.2%) | 2 (5.4%) | 3 (30.0%) | 6 (19.4%) | 9 (24.3%) | 4 (12.1%) | |
| 10-15 years | 32 (21.6%) | 5 (13.5%) | 1 (10.0%) | 4 (12.9%) | 7 (18.9%) | 15 (45.5%) | |
| 15-20 years | 31 (20.9%) | 10 (27.0%) | 2 (20.0%) | 9 (29.0%) | 2 (5.4%) | 8 (24.2%) | |
| >20 years | 38 (25.8%) | 17 (46.0%) | 4 (40.0%) | 4 (12.9%) | 7 (18.9%) | 6 (18.2%) | |
| Number of patients seen in 1 week | | | | | | | 0.003* |
| ≥30-50 | 66 (44.6%) | 21 (56.8%) | 1 (10.0%) | 20 (64.5%) | 8 (21.6%) | 16 (48.5%) | |
| 51-65 | 14 (9.5%) | 3 (8.1%) | - | 1 (3.2%) | 6 (16.2%) | 4 (12.1%) | |
| >65 | 68 (45.9%) | 13 (35.1%) | 9 (90.0%) | 10 (32.3%) | 23 (62.2%) | 13 (39.4%) | |
| Pathologies suffered by patients | | | | | | | |
| Parkinson | 13.3% | 13.0% | 8.3% | 15.6% | 11.7% | 14.8% | 0.296 |
| Stroke | 12.2% | 7.8% | 5.8% | 13.4% | 12.6% | 17.5% | 0.006* |
| Migraine | 35.2% | 30.4% | 25.0% | 38.9% | 43.9% | 30.6% | 0.012* |
| Dementia/Alzheimer | 10.7% | 9.5% | 14.0% | 12.1% | 12.5% | 7.7% | 0.074 |
| Other | 28.5% | 39.2% | 46.9% | 20.0% | 19.3% | 29.4% | <0.001 |

| | TOTAL (n=148) | France (n=37) | Germany (n=10) | Italy (n=31) | Spain (n=37) | UK (n=33) | p value |
|--|------------------|------------------|-------------------|-----------------|-----------------|-----------|---------|
| Type of migraine suffered by the patients | | | | | | | |
| Migraine without aura | 39.9% | 47.3% | 46.5% | 35.3% | 40.7% | 33.1% | 0.003* |
| Migraine with aura | 21.0% | 19.9% | 28.5% | 22.0% | 18.2% | 22.2% | 0.057 |
| Chronic migraine | 26.1% | 21.5% | 17.7% | 26.5% | 27.5% | 31.6% | 0.016* |
| Probable migraine | 10.8% | 12.5% | 7.0% | 8.0% | 14.7% | 8.2% | 0.059 |
| Episodic symptoms that may be associated with migraine | 7.9% | 6.4% | 4.8% | 8.2% | 8.2% | 9.8% | 0.114 |

*p<0.05