

**TableS2.** Tools used to assess treatment in patients with episodic or chronic migraine, by country

	<b>TOTAL (n=148)</b>	<b>France (n=37)</b>	<b>Germany (n=10)</b>	<b>Italy (n=31)</b>	<b>Spain (n=37)</b>	<b>UK (n=33)</b>	<b>p value</b>
<b>First visit – diagnosis</b>							
Clinical interview	147 (99.3%)	37 (100%)	10 (100%)	30 (96.8%)	37 (100%)	33 (100%)	0.434
Patient diary	61 (41.2%)	18 (48.7%)	7 (70%)	13 (41.9%)	16 (43.2%)	7 (21.2%)	0.044*
Validated tools to assess chronic migraine	53 (35.8%)	12 (32.4%)	3 (30%)	15 (48.4%)	14 (37.8%)	9 (27.3%)	0.466
Nothing	1 (0.7%)	-	-	1 (3.2%)	-	-	0.434
<b>Follow up</b>							
Clinical interview	131 (88.5%)	33 (89.2%)	10 (100%)	28 (90.3%)	34 (91.9%)	26 (78.8%)	0.298
Patient diary	127 (85.8%)	34 (91.9%)	8 (80%)	30 (96.8%)	33 (89.2%)	22 (66.7%)	0.005*
Validated tools to assess chronic migraine	54 (36.5%)	9 (24.3%)	2 (20%)	17 (54.8%)	9 (24.3%)	17 (51.5%)	0.009*
Nothing	-	-	-	-	-	-	>0.999
<b>Before starting a preventive treatment</b>							
Clinical interview	140 (94.6%)	36 (97.3%)	10 (100%)	28 (90.3%)	36 (97.3%)	30 (90.9%)	0.461
Patient diary	129 (87.2%)	31 (83.8%)	9 (90%)	30 (96.8%)	32 (86.5%)	27 (81.8%)	0.424
Validated tools to assess chronic migraine	53 (35.8%)	13 (35.1%)	3 (30%)	14 (45.2%)	13 (35.1%)	10 (30.3%)	0.777
Nothing	-	-	-	-	-	-	>0.999
<b>During a preventive treatment</b>							
Clinical interview	135 (91.2%)	34 (91.9%)	10 (100%)	28 (90.3%)	36 (97.3%)	27 (81.8%)	0.174
Patient diary	140 (94.6%)	33 (89.2%)	10 (100%)	29 (93.5%)	36 (97.3%)	32 (97.9%)	0.456
Validated tools to assess chronic migraine	62 (41.9%)	12 (32.4%)	3 (30%)	15 (48.4%)	15 (40.5%)	17 (51.5%)	0.439
Nothing	-	-	-	-	-	-	>0.999

\*p<0.05