

TableS3. Criteria used to assess the effectiveness of a preventive treatment in episodic and chronic migraine, by country.

	Total (n=148)	France (n=37)	Germany (n=10)	Italy (n=31)	Spain (n=37)	UK (n=33)	p value
Frequency of the attacks							0.027*
Always	132 (89.2%)	37 (100%)	10 (100%)	25 (80.6%)	35 (94.6%)	25 (75.8%)	
Usually	15 (10.1%)	-	-	6 (19.4%)	2 (5.4%)	7 (21.2%)	
Sometimes	1 (0.7%)	-	-	-	-	1 (3%)	
Treatment tolerability							0.002*
Always	108 (73.0%)	33 (89.2%)	9 (90.0%)	17 (54.8%)	30 (81.1%)	18 (54.6%)	
Usually	33 (22.3%)	3 (8.1%)	1 (10.0%)	9 (29.1%)	6 (16.2%)	14 (42.4%)	
Sometimes	7 (4.7%)	1 (2.7%)	-	5 (16.1%)	1 (2.7%)	1 (3.0%)	
Intensity of the attacks							0.835
Always	108 (73.0%)	26 (70.3%)	9 (90.0%)	23 (74.2%)	28 (75.7%)	22 (66.7%)	
Usually	33 (22.3%)	8 (21.6%)	1 (10.0%)	7 (22.6%)	7 (18.9%)	10 (30.3%)	
Sometimes	7 (4.7%)	3 (8.1%)	-	1 (3.2%)	2 (5.4%)	1 (3.0%)	
Use of acute migraine medications							0.030*
Always	86 (58.1%)	28 (75.7%)	6 (60.0%)	17 (54.8%)	19 (51.4%)	16 (48.5%)	
Usually	47 (31.8%)	7 (18.9%)	4 (40.0%)	8 (25.8%)	14 (37.8%)	15 (45.5%)	
Sometimes	12 (8.1%)	-	-	6 (19.4%)	4 (10.8%)	2 (6.0%)	
Rarely	3 (2.0%)	2 (5.4%)	-	-	-	-	
Patient's perception of treatment							0.022*
Always	77 (52.0%)	26 (70.3%)	7 (70.0%)	8 (25.8%)	20 (54.1%)	16 (48.5%)	
Usually	49 (33.1%)	8 (21.6%)	3 (30.0%)	16 (51.6%)	7 (18.9%)	15 (45.5%)	
Sometimes	18 (12.2%)	3 (8.1%)	-	5 (16.2%)	8 (21.6%)	2 (6.0%)	
Rarely	3 (2.0%)	-	-	1 (3.2%)	2 (5.4%)	-	
Never	1 (0.7%)	-	-	1 (3.2%)	-	-	

	Total (n=148)	France (n=37)	Germany (n=10)	Italy (n=31)	Spain (n=37)	UK (n=33)	p value
Duration of the attacks							0.403
Always	76 (51.4%)	24 (64.9%)	5 (50.0%)	13 (41.9%)	19 (51.4%)	15 (45.5%)	
Usually	47 (32.4%)	9 (24.3%)	5 (50.0%)	10 (32.3%)	11 (29.7%)	13 (39.4%)	
Sometimes	24 (15.5%)	4 (10.8%)	-	8 (25.8%)	7 (18.9%)	4 (12.1%)	
Rarely	1 (0.7%)	-	-	-	-	1 (3.0%)	
Response to acute migraine medications							0.165
Always	69 (46.6%)	25 (67.6%)	3 (30.0%)	13 (41.9%)	14 (37.9%)	14 (42.4%)	
Usually	50 (33.8%)	10 (27.0%)	4 (40.0%)	11 (35.5%)	13 (35.1%)	12 (36.4%)	
Sometimes	22 (14.9%)	2 (5.4%)	1 (10.0%)	6 (19.4%)	8 (21.6%)	5 (15.2%)	
Rarely	7 (4.7%)	-	2 (20.0%)	1 (3.2%)	2 (5.4%)	2 (6.0%)	

*p<0.05